

RESEARCH OUTPUTS / RÉSULTATS DE RECHERCHE

Innate immune recognition of flagellin limits systemic persistence of *Brucella*

Terwagne, Matthieu; Ferooz, Jonathan; Rolán, Hortensia G.; Sun, Yao Hui; Atluri, Vidya; Xavier, Mariana N.; Franchi, Luigi; Núñez, Gabriel; Legrand, Thomas; Flavell, Richard A.; De Bolle, Xavier; Letesson, Jean-Jacques; Tsolis, Renée M.

Published in:
Cellular microbiology

DOI:
[10.1111/cmi.12088](https://doi.org/10.1111/cmi.12088)

Publication date:
2013

Document Version
Peer reviewed version

[Link to publication](#)

Citation for published version (HARVARD):

Terwagne, M, Ferooz, J, Rolán, HG, Sun, YH, Atluri, V, Xavier, MN, Franchi, L, Núñez, G, Legrand, T, Flavell, RA, De Bolle, X, Letesson, J-J & Tsolis, RM 2013, 'Innate immune recognition of flagellin limits systemic persistence of *Brucella*', *Cellular microbiology*, vol. 15, no. 6, pp. 942-960. <https://doi.org/10.1111/cmi.12088>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Innate immune recognition of flagellin limits systemic persistence of Brucella

Journal:	<i>Cellular Microbiology</i>
Manuscript ID:	Draft
Manuscript Type:	Research article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>TERWAGNE, MATTHIEU; University of Namur, Research Unit in Microorganisms Biology</p> <p>FEROOZ, JONATHAN; University of Namur, Research Unit in Microorganisms Biology</p> <p>Rolan, Hortensia; University of California, Department of Medical Microbiology & Immunology</p> <p>Sun, Yao-Hui; University of California at Davis, Medical Microbiology & Immunology</p> <p>Atluri, Vidya; University of California, Department of Medical Microbiology & Immunology</p> <p>Xavier, Mariana; University of California, Department of Medical Microbiology & Immunology</p> <p>FRANCHI, LUIGI; University of Michigan Medical School, Department of Pathology,</p> <p>Nunez, Gabriel; University of Michigan, Pathology and Comprehensive Cancer Center</p> <p>LEGRAND, THOMAS; University of Namur (FUNDP), Research Unit in Microorganisms Biology; University of Namur, Research Unit in Microorganisms Biology</p> <p>Flavell, Richard; Yale University School of Medicine, Calendar Contact Us Maps & Directions Yale Phonebook YS Department of Immunobiology</p> <p>De Bolle, Xavier; University of Namur (FUNDP), Research Unit in Microorganisms Biology</p> <p>Letesson, Jean-Jacques; University of Namur (FUNDP), Research Unit in Microorganisms Biology</p> <p>Tsolis, Renee; University of California, Medical Microbiology & Immunology</p>
Key Words:	flagellin, Brucella, innate immunity, NLRC4, ASC

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Innate immune recognition of flagellin limits systemic persistence of *Brucella*

2 Matthieu **Terwagne**¹, Jonathan **Ferooz**¹, Hortensia G. **Rolán**², Yao-Hui **Sun**², Vidya **Atluri**²,

3 Mariana N. **Xavier**², Luigi **Franchi**³, Gabriel **Núñez**³, Thomas **Legrand**¹, Richard A.

4 Flavell⁴, Xavier De **Bolle**¹, Jean-Jacques **Letesson**^{1*}, Renée M. **Tsolis**^{2*}

5 ¹URBM, University of Namur (FUNDP), Belgium

6 ²Department of Medical Microbiology & Immunology, University of California, Davis

7 ³Department of Pathology, University of Michigan Medical School, Ann Arbor

8 ⁴ Department of Immunobiology, Yale University School of Medicine, New Haven, CT

9 06520, USA

10 * Both authors contributes equally to this work

11 **Corresponding authors :**

12 Jean-jacques Letesson.

13 Mailing address : URBM, Université de Namur (FUNDP). Rue de Bruxelles 61, 5000 Namur,

14 Belgique.

15 E-mail : jean-jacques.letesson@fundp.ac.be.

16 Tel : (+32) 81 72 44 02.

17 Fax : (+32) 81 72 44 20.

18 Renée M. Tsolis.

19 Mailing address : Medical Microbiology & Immunology, UC Davis, One Shields Avenue,

20 Davis, CA 95616, USA

21 E-mail : rmtsolis@ucdavis.edu

22 Tel : (+1) 530 754 8497

23 Fax : (+1) 530 752 7240

24 **Running title:** *Brucella* flagellin as innate immune signal

25 _____

Abstract

Brucella are facultative intracellular bacteria that cause chronic infections by limiting innate immune recognition. It is currently unknown whether *Brucella* FliC flagellin, the monomeric subunit of flagellar filament, is sensed by the host during infection. Here, we used two mutants of *Brucella melitensis*, either lacking or overexpressing flagellin to show that FliC hinders bacterial replication *in vivo*. The use of cells and mice genetically deficient for different components of inflammasomes suggested that FliC was a target of the cytosolic innate immune receptor NLRC4, and that the cytosolic adaptor ASC was involved in its recognition. Accordingly, we showed that FliC was translocated into the cytosol of infected cells. However, our work also suggested that the lack of TLR5 activity of *Brucella* flagellin and the tight regulation of its synthesis and/or delivery into host cells are both part of the stealthy strategy of *Brucella* towards the innate immune system. Nevertheless, since a flagellin-deficient mutant of *B. melitensis* was found to cause histologically demonstrable injuries in the spleen of infected mice, we suggested that recognition of FliC during infection plays a crucial role in the immunologic standoff between *Brucella* and its host, which is characterized by a persistent infection with limited inflammatory pathology.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

43 **Introduction**

44 The mammalian innate immune system relies on a limited number of pattern recognition
45 receptors (PRRs) to detect microbial-derived molecules during infection and subsequently
46 trigger an appropriate immune response to the invading pathogen. These microbial features
47 are often referred to as PAMPs for pathogen-associated molecular patterns. The PRRs include
48 toll-like receptors (TLRs), which sense PAMPs on the cell surface or in endosomes (Kawai *et al.*
49 *al.*, 2011), and Nod-like receptors (NLRs), which are cytosolic receptors responding to
50 PAMPs and endogenous danger signals (Lamkanfi *et al.*, 2009, Brodsky *et al.*, 2009a). After
51 stimulus recognition, TLRs initiate multiple signalling pathways involved in the innate
52 inflammatory and antimicrobial responses, as well as in the initiation and control of adaptive
53 immune responses (Kawai *et al.*, 2011). In contrast, upon stimulation, several NLRs,
54 including NLRP1 (also known as NALP1), NLRP3 (NALP3 or cryopyrin), and NLRC4 (Ipaf)
55 assemble inflammasomes, which are multiprotein complexes responsible for activation of the
56 inflammatory cysteine protease caspase-1 (Schroder *et al.*, 2010).
57 Bacterial flagellin, the monomeric subunit of flagellar filament, is a PAMP for both systems.
58 Extracellular flagellin is detected by TLR5 (Hayashi *et al.*, 2001) that activates the MyD88-
59 dependent signalling pathway, leading to the nuclear translocation of NF- κ B, and the
60 activation of mitogen activated protein kinases (MAPK), ultimately inducing the secretion of
61 proinflammatory cytokines and chemokines, such as IL-8 (Gewirtz *et al.*, 2001, Eaves-Pyles
62 *et al.*, 2001, Yu *et al.*, 2003). On the other hand, flagellin injected into the cytoplasm of
63 macrophages through bacterial virulence-associated secretion systems is sensed by NLRC4 in
64 association with NAIP5, another member of the NLR family (Kofoed *et al.*, 2011, Zhao *et al.*,
65 2011). Activation of caspase-1 within the NLRC4 inflammasome leads to the maturation and
66 release of biologically active proinflammatory cytokines IL-1 β and IL-18 (van de Veerdonk *et al.*
67 *al.*, 2011). Moreover, this inflammasome can trigger a proinflammatory form of cell death

known as pyroptosis, (Bergsbaken *et al.*, 2009). Finally, it has been shown that NLRC4 plays a role in maintaining a normal endosome-lysosome trafficking of phagocytized bacteria within macrophages (Amer *et al.*, 2006, Akhter *et al.*, 2009). There is evidence that both TLR5 and NLRC4 play a role in controlling *in vivo* infections caused by pathogenic bacteria including *Salmonella enterica* serotype Typhimurium (Feuillet *et al.*, 2006), *Legionella pneumophila* (Hawn *et al.*, 2003) and *Pseudomonas aeruginosa* (Feuillet *et al.*, 2006, Franchi *et al.*, 2012). However, bacterial countermeasures to avoid flagellin recognition by the innate immune system have also been described. *Helicobacter pylori* and *Campylobacter jejuni* escape TLR5 recognition as a result of changes in the amino acid sequence of flagellin (Andersen-Nissen *et al.*, 2005), and it has been suggested that *S. Typhimurium* downregulates *fliC* expression during macrophage infection to avoid a deleterious strong activation of NLRC4 inflammasome (Cummings *et al.*, 2006, Miao *et al.*, 2010a).

Brucella spp. are Gram-negative bacteria that cause brucellosis, a zoonosis of worldwide importance. In the natural reservoir hosts, including wild and domestic animals, these intracellular pathogens cause abortion and infertility. Humans are accidental hosts and *Brucella melitensis* and *B. abortus* are the most frequent cause of human infection (Corbel, 1997). A key characteristic of *Brucella* infection is its chronic nature. Indeed, animals can remain infected for years, and *Brucella* causes a protracted debilitating disease in untreated humans that can result in serious clinical complications (Young, 1995). As a result, brucellosis has an important economic impact on livestock and remains a major public health concern in endemic countries (Pappas *et al.*, 2006).

An important aspect of *Brucella* virulence is its capacity to survive, replicate and persist within infected cells (Atluri *et al.*, 2011). Persistence of *Brucella* within cells relies at least in part on its ability to control the intracellular trafficking of its vacuole in order to avoid lysosomal degradation and to gain access to its replicative niche derived from the

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

93 endoplasmic reticulum (Anderson *et al.*, 1986). Moreover, the success of *Brucella* lies in its
94 stealthy strategy to cope with the innate immune system. First, the structural features of the
95 *Brucella* envelope allow it to avoid sustained recognition by PRRs and subsequent strong
96 inflammatory responses at the onset of infection (Barquero-Calvo *et al.*, 2007). For example,
97 *Brucella* produces a lipopolysaccharide that signals poorly through TLR4, compared to other
98 bacteria (Lapaque *et al.*, 2006, Barquero-Calvo *et al.*, 2007). In addition, *Brucella* can
99 actively control the inflammatory response by producing a protein that interferes with TLR-
100 dependent signalling pathways (Salcedo *et al.*, 2008, Radhakrishnan *et al.*, 2009, Sengupta *et*
101 *al.*, 2009). Along with the lack of cytotoxicity of *Brucella* for highly parasitized host cells, all
102 the above-mentioned features could render it less noticeable by the host innate immune
103 system than other pathogens (Gross *et al.*, 2000, Barquero-Calvo *et al.*, 2007, Salcedo *et al.*,
104 2008). Nonetheless *Brucella* spp. have virulence factors such as a VirB type IV secretion
105 system (T4SS) (O'Callaghan *et al.*, 1999), cyclic β -1,2-glucan (Briones *et al.*, 2001, Arellano-
106 Reynoso *et al.*, 2005) and flagellar genes (Fretin *et al.*, 2005) that are required for *Brucella* to
107 persist within its host. Although our previous studies focused on the flagellum and its role in
108 persistent infection, it is unknown whether *Brucella* flagellin, FliC, is sensed by the host
109 during infection. Here, we combined host and pathogen genetic approaches to assess the
110 potential of *Brucella* flagellin to stimulate innate immune responses.

111

Results

Mice fail to control infection by flagellin-deficient *B. melitensis* mutants.

In a previous study, insertional inactivation of genes located in the three flagellar loci of *B. melitensis* was reported to result in a marked attenuation of its virulence in mice (Fretin *et al.*, 2005). At that time, it was assumed that, as described in enterobacteriaceae, the *fliC* gene was not expressed in mutants of genes encoding basal flagellar structures. However, we recently demonstrated that the flagellar expression hierarchy of *Brucella* is not conventional, since the flagellin subunit is still produced in mutants deficient in the hook or basal body (Ferooz *et al.*, 2011). To evaluate the specific impact of the absence of FliC flagellin on the virulence of *B. melitensis*, non-polar mutants of *fliC* ($\Delta fliC$) and *flbT* ($\Delta flbT$) (Ferooz *et al.*, 2011) were used to infect murine macrophages and BALB/c mice. The FlbT regulator of *B. melitensis* is specifically required for the production of FliC, most likely by allowing translation of the *fliC* mRNA (Ferooz *et al.*, 2011). Accordingly, flagellin was detected neither in the $\Delta fliC$ nor in the $\Delta flbT$ strain harvested at the early exponential phase of growth, whereas the protein is produced by the isogenic wt strain (Fig. 1A).

We first compared the intracellular growth of *B. melitensis* $\Delta fliC$ and $\Delta flbT$ to that of wt bacteria in RAW264.7 murine macrophages. No difference in colony forming units (CFUs) was detected over a 48-h time course (Fig. 1B). Similar results were obtained in HeLa cells (data not shown). Consistent with a normal multiplication in endoplasmic reticulum-derived vacuoles, both $\Delta fliC$ mutant and its isogenic parental strain were found to replicate within calnexin-positive compartments of HeLa cells at 24h p.i (data not shown).

Despite the absence of an obvious role for *Brucella* flagellar genes in cellular models of infection, several reports have shown that they are required for the establishment of a persistent infection *in vivo* (Fretin *et al.*, 2005, Zygmunt *et al.*, 2006). To re-evaluate the role of flagellar proteins *in vivo*, BALB/c mice were infected via the intraperitoneal route with *B.*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

melitensis 16M $\Delta fliC$, $\Delta flbT$ and $\Delta fliF$ non-polar mutants. None of the mutants was significantly attenuated 5 days p.i., as compared with the parental strain (Fig. 1C). Moreover, we could confirm that the basal body protein FliF is required for full virulence. Indeed, the $\Delta fliF$ mutant was attenuated at 3 and 4 weeks p.i. (Fig. 1C). In contrast, the virulence of the $\Delta fliC$ strain was exacerbated when compared to its isogenic parental strain, as $\Delta fliC$ -infected mice presented a higher bacterial load in the spleen from 12 days until 60 days p.i. (Fig. 1C). A higher bacterial count was also observed at the same times in the livers of mice infected with the $\Delta fliC$ mutant (data not shown). Similarly, an enhanced persistence of the $\Delta fliC$ strain in the spleens of the resistant C57BL/6 mice has also been observed (data not shown). The use of a low-copy plasmid carrying *fliC* gene along with its predicted flanking regulatory sequences, which restores regulated production of flagellin in the $\Delta fliC$ strain (Fig. 1A), allowed partial complementation of the phenotype of the newly constructed $\Delta fliC$ mutant at 28 days p.i. and full complementation at 60 days p.i. (Fig. 1C). Moreover, we could show that the $\Delta flbT$ mutant had similar infection kinetics than the $\Delta fliC$ strain in the spleen of BALB/c mice (Fig. 1C). This further supports the fact that the apparent inability of the host to control bacterial infection is specifically due to the lack of flagellin production by *Brucella*.

Mice infected with *B. melitensis* $\Delta fliC$ mutant exhibit severe splenic pathology.

Brucella is known to induce splenomegaly in infected hosts. During the course of a *B. melitensis* 16M infection in BALB/c mice, the spleen weight increases and peaks around 0.4 gr (4-fold the spleen weight of an uninfected mice) at 12 days p.i. Afterwards, the spleen weight decreases but remains twice the normal value until the end of the experiment (Fig. 2A). In contrast, we found that the splenomegaly of mice infected with flagellin-deficient mutants, while displaying kinetics similar to those of the wt infection during the first 12 days, continued to increase until 28 days p.i. and reached a plateau of almost 5 or 6 times the

1
2
3 162 normal spleen weight by the end of the experiment (Fig. 2A for $\Delta fliC$, data not shown for
4
5 163 $\Delta flbT$). A similar exacerbation of splenomegaly was also observed in C57BL/6 mice at 21
6
7 164 days p.i with the $\Delta fliC$ mutant (data not shown). This was in accordance with the enhanced
8
9 165 persistence of the flagellin-deficient mutants in mice (Fig. 1C).

10
11 166 We further examined the splenic histopathology of BALB/c mice infected for 28 days with wt
12
13 167 or $\Delta fliC$ *B. melitensis* strain. At this time, mice infected with the $\Delta fliC$ strain showed a
14
15 168 markedly exacerbated splenic inflammation characterized by increased vasodilation,
16
17 169 thrombosis, neutrophil infiltration and granuloma formation (Fig 2B and 2C). In contrast,
18
19 170 mice infected for 28 days with wt *B. melitensis* had nearly normal splenic morphology, as
20
21 171 compared with non-infected mice.
22
23
24
25
26

27 173 **Ectopic production of flagellin attenuates the virulence of *B. melitensis* in vivo.**

28
29 174 Mice apparently fail to control infection caused by *B. melitensis* 16M $\Delta fliC$ or $\Delta flbT$ at late
30
31 175 time points. This observation suggests that production of flagellin by *Brucella* somehow
32
33 176 influences the course of infection. To further test this hypothesis, we engineered a *B.*
34
35 177 *melitensis* 16M strain, designated *BruFliC^{ON}*, that constitutively expresses a plasmid-encoded
36
37 178 copy of *fliC* from *Escherichia coli* *Plac*. Western blot analysis confirmed that, while
38
39 179 production of flagellin by wt bacteria is only detectable at the early exponential phase of
40
41 180 growth, *BruFliC^{ON}* produced higher levels of flagellin throughout *in vitro* growth (Fig. 3A).

42
43 181 Ectopic production of flagellin did not impair the invasion and replication abilities of *Brucella*
44
45 182 in macrophages *in vitro* (Fig. 3B). However, we found that the *BruFliC^{ON}* strain was
46
47 183 attenuated *in vivo* compared with wt *B. melitensis* 16M. While no difference in splenic
48
49 184 bacterial load was observed between the two strains at 5 days post infection of BALB/c mice,
50
51 185 0.5 to 1 log fewer CFU of *BruFliC^{ON}* bacteria were recovered at 12, 21 and 28 days p.i. (Fig.
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

3C). Reduced colonization of *BruFliC*^{ON} was also observed in the liver of infected BALB/c, and similar results were also obtained with C57BL/6 mice (data not shown).

***Brucella* flagellin lacks TLR5 agonist activity**

The altered virulence of the $\Delta fliC$ and *BruFliC*^{ON} mutants led us to hypothesize that *Brucella* flagellin is detected by the host in order to mount a protective immune response. To ascertain whether innate immune sensing of flagellin contributes to enhanced control of systemic *Brucella* infection, we first determined whether *Brucella* flagellin possesses agonist activity for TLR5. To this end, epitope-tagged FliC flagellins from *Brucella* (*BruFliC*-FLAG) or *S. enterica* serotype Typhimurium (*S. Typhimurium*; *StFliC*-FLAG) were expressed in an *S. Typhimurium* *fliCfljB* mutant (EHW26) lacking endogenous flagellin expression. Immunoblotting with the anti-FLAG antibody demonstrated that both *BruFliC*-FLAG and *StFliC*-FLAG were secreted to the supernatant in similar amounts (Fig. 4A). Addition of the C-terminal FLAG tag to *StFliC* prevents its assembly into flagellar filaments, thereby allowing for a direct comparison of effects of flagellin monomers in the absence of a confounding effect on motility, since strains expressing either *StFliC*-FLAG or *BruFliC*-FLAG were aflagellate and non-motile (data not shown).

Culture supernatants of *S. Typhimurium* *fliCfljB* expressing recombinant flagellins were used to treat two TLR5-expressing cell lines: the colonic epithelial cell line T84 and HEK293/hTLR5 (Fig. 4B and 4C). Both cell lines secreted interleukin 8 (IL-8) on infection with strains expressing native or FLAG-tagged *StFliC*, demonstrating that addition of the epitope tag to the C terminus of flagellin did not affect its TLR5 agonist activity. Stimulation of IL-8 secretion was dependent on flagellin in both cell lines, since culture supernatants from the *fliCfljB* mutant elicited little (Fig. 4C) or no (Fig. 4B) IL-8. In contrast to *StFliC*-FLAG, expression of *BruFliC*-FLAG did not elicit IL-8 secretion above the level of the *fliCfljB*

mutant. Similar results were obtained when T84 or HEK-293/hTLR5 cells were infected with *S. Typhimurium* strains expressing recombinant flagellins (data not shown). The response to *BruFliC* did not appear to be delayed, since extending the time of the assay to 24h did not allow detection of a response comparable to that elicited by StFliC-FLAG. As a second readout for TLR5 signaling, we assayed activation of mitogen-activated protein kinases (MAPK) p38 and ERK by treatment with purified, GST-tagged flagellins. Phosphorylation of both p38 and ERK was induced to a greater extent by GST-StFliC than by GST-*BruFliC*, and notably no increase in phosphorylation of ERK could be detected after treatment with GST-*BruFliC* (Fig. 4D). Taken together, these results demonstrate that compared to *S. Typhimurium* flagellin, the ability of *Brucella* flagellin to stimulate TLR5 signaling is greatly reduced.

Cytosolic sensing pathways detect *Brucella* flagellin during infection of macrophages

In addition to TLR5, flagellin that enters the cytosol of host macrophages can be sensed by the NLRC4/NAIP5 pathway (Kofoed *et al.*, 2011, Zhao *et al.*, 2011). To determine whether cytosolic pathways could detect flagellin during *Brucella* infection, we first used the TEM-1 β -lactamase assay to detect translocation of flagellin into the cytosol of *B. abortus*-infected J774 macrophage-like cells. For these experiments, J774 cells were infected with *B. abortus* 2308 expressing either a C-terminally tagged copy of *Brucella* flagellin or an irrelevant protein (GST), from a multi-copy plasmid (pFLAG-TEM1; Sun *et al.*, 2007). While cells infected with *B. abortus* expressing GST::Flag-TEM-1 showed no cytosolic β -lactamase activity (no β -lactamase-positive cells in 4 experiments), 0.94% (range: 0.3-2.1%) of cells infected with *B. abortus* expressing the flagellin fusion protein were β -lactamase positive, suggesting potential access of low amounts of flagellin to the cytosol of *Brucella*-infected cells. Next, we determined whether, in primary macrophages, cytosolic flagellin could

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

stimulate innate immune responses. To this end, we compared the ability of *B. melitensis* and its isogenic $\Delta fliC$ mutant to elicit IL-1 β secretion from primary bone marrow-derived macrophages (BMDM). Compared to *B. melitensis* wt, the $\Delta fliC$ mutant elicited significantly reduced IL-1 β secretion (Fig. 5A). This reduction was not the result of differing numbers of intracellular bacteria of the $\Delta fliC$ mutant, since both the $\Delta fliC$ mutant and wt *B. melitensis* were present in the same numbers (data not shown). This partial reduction in IL-1 β secretion suggests that recognition of flagellin contributes to activation of the caspase-1 inflammasome. The mechanism of cytosolic flagellin sensing in the context of intracellular infection was further investigated using the *B. melitensis* FliC^{ON} strain, which expresses flagellin constitutively. This strain, as well as a control carrying the empty plasmid pBBR1MCS, was used to infect immortalized BMDM from mice deficient in NLRC4 (Fig. 5B). Constitutive expression of FliC did not affect the ability of *B. melitensis* to survive intracellularly (Fig. 3B and data not shown). The BruFliC^{ON} strain elicited significantly more IL-1 β secretion than the control strain, confirming data shown in Fig 5A. While these results suggested that under conditions of flagellin expression, flagellin can be sensed by cytosolic PRRs that lead to activation of caspase-1 and secretion of IL-1 β , NLRC4 was not required for flagellin-dependent stimulation of IL-1 β secretion by BMDM *in vitro*.

***Brucella* flagellin elicits IL-1 β secretion by a mechanism that is distinct from the NLRC4/NAIP5 pathway**

Since *B. melitensis* is known to inhibit innate immune signalling (Salcedo *et al.*, 2008, Radhakrishnan *et al.*, 2009, Sengupta *et al.*, 2009) we determined whether purified flagellin, in the absence of other *Brucella* factors, would signal similarly to flagellin expressed during cellular infection. For this purpose, purified GST-BruFliC and GST-StFliC were introduced into the cytosol of BMDM using the cationic lipid DOTAP (Franchi *et al.*, 2006). Both

261 *Bru*FliC and StFliC elicited dose-dependent secretion of IL-1 β from BMDM from C57BL/6
 262 mice when introduced into the cytosol using DOTAP (Fig. 6A). Neither GST, DOTAP alone,
 263 nor recombinant flagellins in the absence of DOTAP elicited any secretion of IL-1 β (Fig. 6A
 264 and data not shown). Comparison of IL-1 β secreted in response to equal amounts of StFliC or
 265 *Bru*FliC suggested that the proinflammatory activity of StFliC was slightly higher than that of
 266 *Bru*FliC (Fig. 6A). Secretion of IL-1 β in response to *S. Typhimurium* FliC was dependent on
 267 NLRC4 and only partially dependent on the adaptor protein ASC (apoptosis-associated speck-
 268 like protein), as reported previously (Broz *et al.*, 2010). In contrast, *Bru*FliC elicited IL-1 β
 269 secretion that required ASC, but was independent of NLRC4, at least in cultured BMDM
 270 (Fig. 6B and Fig. 6C). These results suggested that in BMDM, *Brucella* flagellin was sensed
 271 by a cytosolic mechanism that differs from the NLRC4/NAIP5-dependent response to *S.*
 272 *Typhimurium* FliC (Kofoed *et al.*, 2011, Zhao *et al.*, 2011).

273
 274 **The cytosolic flagellin-detection pathway is implicated in the control of *B. melitensis***
 275 **infection *in vivo*.**

276 To evaluate the potential impact of caspase-1 inflammasomes on the control of *Brucella*
 277 infection *in vivo*, we infected *Nlrc4*^{-/-} and *Casp1*^{-/-} C57BL/6 mice with *B. melitensis* 16M.
 278 Splenic bacterial count was examined 21 days p.i., a time at which wt mice manage to
 279 effectively control infection caused by flagellin-producing *Brucella* strains (Fig. 1C and 3C).
 280 At this time, we observed that NLRC4 (Fig. 7A) and caspase-1 (Fig. 7B) deficiency
 281 moderately but significantly affected the resistance of mice to infection. This suggests that the
 282 NLRC4-caspase-1 axis is required for the host to control *B. melitensis* 16M infection,
 283 possibly through recognition of cytosolic flagellin. To further test this hypothesis, the
 284 *Bru*FliC^{ON} strain was used to infect *Nlrc4*^{-/-} and *Casp1*^{-/-} mice. As shown previously, virulence
 285 of this strain is attenuated compared to wt *B. melitensis* 16M, as the spleen of *Bru*FliC^{ON}-

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

infected C57BL/6 mice contained less CFUs than these infected by the wt strain (Fig. 7). Interestingly, this virulence defect was rescued in mice deficient for the cytosolic flagellin sensor NLRC4 (Fig. 7A) or the downstream caspase-1 (Fig. 7B). These data indicate that, in contrast to what has been observed *in vitro* (Fig. 6), *Brucella* flagellin can activate the NLRC4 inflammasome *in vivo*. Nevertheless, although *Nlrc4*^{-/-} and *Casp1*^{-/-} mice infected with wt *B. melitensis* 16M had significantly higher splenic bacterial counts than those of wt mice, it remained significantly lower than those of mice infected with the Δ *fliC* mutant (Fig. 7). This suggests that both inflammasome-dependent and inflammasome-independent control of infection operates downstream detection of *Brucella* flagellin *in vivo*.

***B. melitensis* Δ *fliC* mutant fails to elicit early granuloma formation in the spleen of infected mice.**

Chronic granulomatous inflammation in the spleen of natural hosts, humans and mice is the hallmark of *Brucella* infection (Spink *et al.*, 1949, Enright *et al.*, 1990). Recently, we revealed the pivotal role of early splenic granuloma formation in the ability of mice to control bacterial dissemination (Copin *et al.*, 2012). Here, we used a rabbit polyclonal serum raised against *B. melitensis* (anti-*Bru*) with the aim to compare the distribution of putative infected cells in the spleen of BALB/c mice inoculated with *B. melitensis* 16M wt or Δ *fliC* strain. 5 days after infection with *B. melitensis* 16M wt, clusters of cells stained with anti-*Bru* (*Bru*-positive cells) were found equally in white pulp and red pulp area of the spleen (Fig. 8). These clusters consisted primarily of CD11b⁺ cells, suggesting that they corresponded to the granuloma previously described (Copin *et al.*, 2012). Strikingly, at the same time, the number of *Bru*-positive cells clusters counted in splenic sections of Δ *fliC*-infected mice was reduced (Fig. 8). This apparent defect in early splenic granuloma formation suggests the importance of flagellin sensing by the host for the orchestration of this typical tissue response to *Brucella* infection.

311

312 **Discussion**

313 Intracellular survival and immune evasion both contribute to persistence of *Brucella* in the
314 host (Atluri *et al.*, 2011). Recent studies have shown that *Brucella* uses passive as well as
315 active mechanisms to evade detection by TLRs of the innate immune system (Lapaque *et al.*,
316 2006, Barquero-Calvo *et al.*, 2007, Salcedo *et al.*, 2008, Radhakrishnan *et al.*, 2009, Sengupta
317 *et al.*, 2009). Accordingly, the inflammatory response induced at the onset of *Brucella*
318 infection is lower than observed with pyogenic infections such as salmonellosis (Barquero-
319 Calvo *et al.*, 2007). Actually, brucellae are not entirely invisible to the immune system, which
320 can still detect them and shape a Th1 response to control infection (Murphy *et al.*, 2001,
321 Copin *et al.*, 2007). However, the host immune response is not sufficient to eliminate bacteria,
322 resulting in a chronic state of infection characterized by a balance between pathogen virulence
323 and host resistance. The impact of *Brucella* flagellin on infection had not been reported yet.
324 The data presented here suggest that flagellin plays a crucial role in the interplay between
325 *Brucella* and its host, as its detection by the innate immune system is required for the control
326 of infection *in vivo*, although some characteristics of *Brucella* flagellin would contribute to
327 the stealthy strategy of this pathogen.

328

329 The use of two mutants of *B. melitensis* 16M that either overproduce or lack the FliC flagellin
330 has shown that this protein hinders bacterial replication *in vivo*. Indeed, a strain engineered to
331 ectopically produce flagellin (*BruFliC^{ON}*) was attenuated in mice, whereas deletion of *fliC*
332 (Δ *fliC*) enhanced persistence of *B. melitensis* 16M in these conditions. Our *in vivo* data are
333 consistent with studies reporting exacerbated infections caused by a flagellin deficient mutant
334 of *Salmonella enterica* serovar Typhimurium (Vijay-Kumar *et al.*, 2006), *Legionella*
335 *pneumophila* (Molofsky *et al.*, 2006) or *Pseudomonas syringae* pv. *Tabaci* (Li *et al.*, 2005),

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

as well as virulence attenuation due to flagellin overproduction by *S. Typhimurium* (Salazar-Gonzalez *et al.*, 2007, Miao *et al.*, 2010a) and *Listeria monocytogenes* (Grundling *et al.*, 2004). These findings also suggest that *Brucella* flagellin is an important immune target during infection, and our work provides first insights into the mechanisms involved.

TLR5 and the NLRC4/NAIP5 complex are the only proteins currently known as innate immune sensors of extracellular and cytoplasmic bacterial flagellin, respectively (Miao *et al.*, 2007).

In agreement with a recent paper quoting that purified *Brucella* flagellin does not induce expression of interferon-inducible resistance proteins (IRGs) in murine macrophages (Lapaque *et al.*, 2009), the data reported in this paper allow us to conclude that *Brucella* flagellin is not a TLR5 agonist. This is consistent with its atypical sequence as it lacks the amino acid residues required to stimulate this PRR (Andersen-Nissen *et al.*, 2005). Thus, we propose that *Brucella* evades TLR5-mediated detection, and that it could be part of its stealthy strategy to avoid activation of the innate immune system during the onset of infection.

Cytosolic flagellin activates a complex comprising the NLR family proteins NLRC4 and NAIP5 (Franchi *et al.*, 2006, Miao *et al.*, 2006, Kofoed *et al.*, 2011, Zhao *et al.*, 2011). This complex senses a highly conserved region of the C terminal part of the flagellin critical for flagellum filament assembly (Yonekura *et al.*, 2003), but that is required neither for flagellin translocation into the host cell cytosol nor for TLR5 activation (Lightfield *et al.*, 2008). The C-terminal 35 amino acid residues are conserved in *Brucella* FliC flagellin, as they share respectively 46% and 40% identity with *L. pneumophila* FlaA and *S. Typhimurium* FliC, both known to activate NLRC4 (Franchi *et al.*, 2006, Zamboni *et al.*, 2006) and sharing themselves 60% identity. Recently, it has been proposed that the minimal motif of flagellin sensed by NLRC4 comprises the highly conserved last C-terminal residues VLSLL found in *L.*

1
2
3 361 *pneumophila* FlaA and *S. Typhimurium* FliC (Lightfield *et al.*, 2008, Miao *et al.*, 2010b).
4
5 362 This motif is semi-conserved in *Brucella* flagellin that bears an ILSFR motif.
6
7 363 Our results suggest that, similar to what is seen with *L. pneumophila* infection (Amer *et al.*,
8
9 364 2006, Case *et al.*, 2009) the NLRC4-caspase-1 axis is involved in the control of *B. melitensis*
10
11 365 16M *in vivo* (Fig. 7). However, the absence of NLRC4 or caspase-1 stimulation in mice
12
13 366 infected with the flagellin-deficient *B. melitensis* 16M Δ *fliC* or Δ *flbT* mutants cannot by itself
14
15 367 account for the inability of the host to control infection. Indeed, the relative difference of
16
17 368 virulence between *B. melitensis* 16M wt and Δ *fliC* strains were only partially reduced in
18
19 369 *Nlrc4*^{-/-} and *Casp1*^{-/-} mice (Fig. 7), indicating involvement of both NLRC4/caspase-1-
20
21 370 dependent and independent mechanisms in the control of *Brucella* downstream flagellin
22
23 371 recognition. This contrasts with what is observed after intratracheal infection of mice with *L.*
24
25 372 *pneumophila*. Indeed, in this case, the number of *flaA* mutants and wt bacteria in the lungs of
26
27 373 *Nlrc4*^{-/-} and *Casp1*^{-/-} is similar (Amer *et al.*, 2006, Case *et al.*, 2009). Therefore, it suggests
28
29 374 that *Brucella* flagellin is an immune target not only for the cytosolic sensor NLRC4 *in vivo*.
30
31 375 Actually, the observation that the *BruFliC*^{ON} strain is still attenuated (a reproducible 0.5 log
32
33 376 decreased CFUs in the spleen) compared to *B. melitensis* 16M wt in *Nlrc4*^{-/-} and *Casp1*^{-/-} mice
34
35 377 21 days p.i. is consistent with the hypothesis that *Brucella* flagellin stimulates another
36
37 378 immune pathway in addition to the NLRC4/caspase-1 axis. The ASC-dependent signalling
38
39 379 suggested by our *ex-vivo* data (Fig. 6) could be this additional pathway. This would be similar
40
41 380 to what has been described for *L. pneumophila* that triggers an ASC-dependent activation of
42
43 381 caspase-1 in macrophages, in addition to the NLRC4-dependent activation triggered by
44
45 382 cytosolic flagellin (Case *et al.*, 2009). Activation of these innate immune pathways by
46
47 383 flagellin would play a role in limiting replication of *Brucella in vivo*. However, the immune
48
49 384 effector mechanisms involved remain to be uncovered. Processing of the proinflammatory
50
51 385 cytokines pro-IL-1 β and pro-IL-18 (Raupach *et al.*, 2006, Dinarello, 2009), pyroptosis
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

(Bergsbaken *et al.*, 2009, Miao *et al.*, 2010a) and control of phagosome maturation (Amer *et al.*, 2006, Akhter *et al.*, 2009) that can all result from caspase-1 activation are important processes for innate immunity against bacterial pathogens (Brodsky *et al.*, 2009b).

Besides its impact on the innate immune system, it is known that bacterial flagellin is also a target of the adaptive immune response (Salazar-Gonzalez *et al.*, 2005). However, whether the adaptive immune system responds to MHC class II-presented flagellin peptides during infection by *Brucella* is currently not known.

While searching for immune effector mechanisms triggered by flagellin detection and involved in the control of *Brucella* replication in mice, we found that the $\Delta fliC$ mutant fails to elicit early granulomatous response in the spleen of mice infected for 5 days, a time at which the mutant is found at a similar level as the wt strain (Fig. 8). Thus, we suggest that detection of flagellin by the host would play a role in early granuloma development during brucellosis. Although the granulomatous response was stronger at 28 days p.i. (Fig. 2B), when the $\Delta fliC$ strain colonized spleens at higher extent than wt, an early alteration in this response could contribute to the apparent failure of mice to control infections caused by the flagellin-deficient mutants of *B. melitensis* 16M. Indeed, granulomatous inflammation is the typical tissue response to *Brucella* infection in both mice and humans (Spink *et al.*, 1949, Hunt *et al.*, 1967, Enright *et al.*, 1990), and a recent study has demonstrated the crucial role of early formation of splenic granuloma in the control of *B. melitensis* 16M (Copin *et al.*, 2012). Whether granuloma formation during infection by *Brucella* depends on ASC, NLRC4 and/or caspase-1 is currently unknown. Up to now, a role for the NLRC4 inflammasome in such a response has never been reported. However, it was recently shown that granuloma formation in chronic *M. tuberculosis* infection is dependent on ASC, whereas it does not require caspase-1 (McElvania Tekippe *et al.*, 2010).

411

412 *S. Typhimurium* translocates flagellin from its containing-vacuole into the cytosol of infected

413 cells by a SPI1-T3SS-dependent but flagellar secretory apparatus-independent process (Sun *et*

414 *al.*, 2007). Similarly, a Dot/Icm T4SS-mediated flagellin translocation has been suggested in

415 the case of *L. pneumophila* (Ren *et al.*, 2006, Molofsky *et al.*, 2006). Here, we show that

416 *Brucella* flagellin is also translocated into the host cell cytosol. Interestingly, flagellin

417 translocation was not seen when a *virB2* mutant was used to infect macrophages (data not

418 shown), suggesting that VirB T4SS may play a role in flagellin translocation. Interestingly, a

419 requirement for the T4SS to elicit splenic microgranuloma formation has been proposed

420 (Rolan *et al.*, 2009). According to our results, it could be envisioned that the VirB T4SS of

421 *Brucella* elicits a granulomatous response by translocating flagellin. However, since the T4SS

422 is also essential for *Brucella* to reach its replicative niche (Celli *et al.*, 2003), additional

423 studies would be necessary to determine whether the role of the T4SS in release of flagellin to

424 the host cytosol is direct or indirect. The TEM1 β -lactamase reporter assay has been

425 previously used to demonstrate translocation of *S. Typhimurium* flagellin into the cytosol of

426 infected macrophages (Sun *et al.*, 2007). We observed that the amount of flagellin

427 translocated into cells by *Brucella* is far less than by *Salmonella*. While flagellin could be

428 detected in the cytosol of 77.5% of macrophages infected for 4h with *S. Typhimurium* (Sun *et*

429 *al.*, 2007), less than 1% of cells were positive 16h after infection with *B. abortus*. Therefore,

430 although the intrinsic ability of *Brucella* and *Salmonella* flagellin to induce IL-1 β secretion

431 from BMDM appeared to be similar (Fig. 6), *Brucella* might evade activation of a robust

432 innate immune response from cytosolic PRRs by controlling the production and/or delivery of

433 flagellin into the host cell. Accordingly, we could show that the attenuation of the *BruFliC*^{ON}

434 strain that ectopically produces flagellin is due to a strong NLRC4 inflammasome activation

435 *in vivo* (Fig. 7). Thus, we propose that the tight regulation of flagellin synthesis and/or

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

delivery during infection is part of its stealthy strategy. This has also been suggested for *S.* Typhimurium, which downregulates the expression of *fliC* during macrophage infection (Cummings *et al.*, 2006).

In conclusion, we propose that flagellin is an important molecular actor of the interplay between *Brucella* and its host. Although flagellin escapes detection by TLR5 and *Brucella* controls its production and/or delivery to the infected host cell cytosol, its detection by cytosolic PRRs initiates a response that results in an immunologic standoff between *Brucella* and its host, leading to a persistent infection with limited inflammatory pathology. The increased bacterial tissue loads and destructive pathology, seen with the flagellin-deficient mutant demonstrates that innate and possibly also adaptive, recognition of flagellin is a process that is important to the chronic and stealthy nature of *Brucella* infection. As such, flagellin could be considered as a “host protective factor” (Shames *et al.*, 2010) in the context of brucellosis.

Experimental procedures

Bacteria and growth conditions

Bacterial strains and plasmids are listed in Table 1. Cultures of *Brucella* strains were freshly inoculated from frozen stock onto 2YT medium (10% yeast extract, 10 g liter⁻¹ tryptone, 5 g liter⁻¹ NaCl) plates before subculturing aerobically at 37°C in 2YT broth supplemented with appropriate antibiotics. LB broth was used for *Escherichia coli* and *Salmonella enterica* serotype Typhimurium (*S.* Typhimurium) cultures. Antibiotics were used at the following

concentrations: carbenicillin, 100 mg/liter; chloramphenicol, 30 mg/liter; kanamycin, 60 mg/liter; or nalidixic acid, 50 mg/liter.

Molecular techniques

DNA manipulations were performed according to standard techniques (Ausubel *et al.*, 1991). Primers used are listed in Table 2.

Generation of the complementation vector pRH001-*fliC*: *fliC* coding sequence (cds) and its predicted upstream and downstream regulatory sequences were amplified by PCR using the *PfliC* and *tfliC* primers pair. The PCR product (*PfliC-fliC-tfliC*) was then cloned into the *EcorRV* site of pGEM. In a second step, this fragment was excised using *BamHI* and *XbaI*, and inserted into the corresponding sites of pMR10*cat* (R. Roberts, unpublished) in the opposite orientation to the *Plac*.

Generation of the *B. melitensis* 16M *FliC*^{ON} strain: The *fliC* overexpression vector pBBR1-*fliC* was obtained as follows: first, the constitutive promoter of the lac operon *Plac* was amplified by PCR using the *Plac* and *fliC-Plac* primers pair. In the resulting PCR product, *Plac* is flanked by translation stop codons in all three reading frame in 5' and by the 21st *fliC* coding sequence (cds) base pairs in 3'. *fliC* cds was amplified by PCR using the *BmfliC-F* and *BmfliC-R* primers. A third PCR using the *Plac* and *BmfliC-R* primers was used to ligate the two PCR products by cohesive ends. Stop codons and close fusion of *fliC* cds to *Plac* without any linker ensure the production of *FliC* flagellin that does not bear additional N-terminal amino acid residues. The PCR product (*XbaI-Plac-fliC-BamHI*) was then cloned into the *EcorRV* site of pGEM. In a last step, this fragment was excised using *XbaI* and *BamHI*, and inserted into the corresponding sites of pBBR1 MCS-I (Kovach *et al.*, 1994) in the opposite orientation to the endogenous *Plac*. This gave rise to pBBR1-*Plac-fliC*. This final

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

construction was transformed into *E. coli* strain S17-1 (Simon *et al.*, 1983), and introduced into *B. melitensis* 16M by conjugation.

Generation of C-terminally FLAG-tagged flagellins: A derivative of the broad host range plasmid pBBR1MCS (pBBR1-FLAG) was first generated by ligating a fragment containing “*SphI*-promoter-*NdeI*-*SalI*-3x-Flag-STOP-*PstI*-*SacI*” into pBBR1MCS4 treated with *SphI* and *SacI*. The *S. Typhimurium fliC* gene was amplified using primers StFliC-F and StFliC-R, and the resulting amplicon was ligated into *NdeI* and *SalI*-digested pBBR1-FLAG to yield plasmid pYHS1116, encoding StFliC-FLAG. The *B. abortus fliC* gene was amplified using primers BaFliC-F and BaFliC-R and cloned in the same way to generate pYHS1073, encoding BaFliC-FLAG. In both constructs, expression of the recombinant proteins was controlled by a previously described constitutive *Brucella* promoter, BMEII0193 (Eskra *et al.*, 2001). The constructs were confirmed by DNA sequencing across the junction fragments. Plasmids pYHS1116 (StFliC-FLAG) and pYHS1073 (BaFliC-FLAG) were introduced into a *Salmonella fliC fljB* mutant (EHW26, (Raffatellu *et al.*, 2005)) by electroporation. The *B. abortus* and *B. melitensis* FliC proteins are identical except for a substitution of Ala156 to Thr in *B. abortus*.

Generation of fusions to TEM-1 β -lactamase: To express BaFliC fused with TEM1, *B. abortus fliC* was amplified by using the primer pair BaFliC-F and BaFliC-R. The amplicon was cloned into pCR2.1, then subsequently digested with *NdeI* and *PstI*, and ligated pFlagTEM1 (Raffatellu *et al.*, 2005) digested with the same enzymes to yield pBaFliCTEM1. The expression of BaFliC::TEM1 in pBaFliCTEM1 is under the control of inducible Trc promoter. Constructs expressing StFliC::TEM1 were described previously (Sun *et al.*, 2007).

Generation of GST-flagellin fusion proteins: For construction of plasmids expressing GST fused at the N-terminus of flagellins, flagellin genes were amplified to delete predicted N-terminal secretion domains. The *fliC* gene from *S. Typhimurium* was amplified without its

first 332 nucleotides using primer pair of StFliC-F2 and StFliC-STOP-R. Similarly *B. abortus* *fliC* lacking its first 87 nucleotides was amplified using primer pair of BaFliC-F2 and BaFliC-R2. Both amplicons were cloned in pCR2.1, excised as *Bam*HI/*Sal*I fragments, and ligated to *Bam*HI/*Sal*I –digested pGEX-4T-1. The cloning junctions were confirmed by DNA sequence analysis, and the resulting constructs, pGEX-StFliC and pGEX-BaFliC, were transformed into *E. coli* BL-21. Expression of GST::StFliC and GST::BaFliC was induced by IPTG, and the recombinant flagellins were purified using Glutathione-Sepharose 4B (GE Healthcare). Protein concentration was measured with DC protein assay (BioRad).

Construction of plasmids expressing native *S. Typhimurium* and *B. abortus* flagellins: The pSC101 *ori*-based low copy number plasmid pWSK29 (Genbank AF016889) was digested with PvuII and BsaAI to remove the *lac* promoter, *lacZ* fragment, and most of the *f1 ori*, as represented by a 4.7kb fragment product (Figure 2a). The 4.7kb plasmid fragment was gel purified then treated with Antarctic phosphatase (New England Biolabs, NEB). Primers 102 and 103 were used to PCR amplify the -134 to -6 region of *S. Typhimurium* LT2, with primer 103 adding an XbaI site which serves to replace the -5 to -1 region of *fliC* and overlaps the translation start site. Primer 103 also contains a multiple cloning site (MCS), adding unique restriction sites (in the context of PvuII/BsaAI-digested pWSK29) downstream of the XbaI site. Both primer 102 and 103 contain PvuII sites at their 5' ends, so the resulting PCR product was cleaved with PvuII, gel purified, and blunt ligated to the pWSK29 PvuII/BsaAI fragment with Quick T4 DNA Ligase (NEB) and heat shocked into *E. coli* DH5 α . Clones were screened for the loss of an EcoRV site (proper PvuII site ligated), gain of a single BstBI site (one promoter region insertion) and the orientation with the *fliC* promoter facing away from the pSC101 *ori* (the same directionality as the *lac* promoter in pWSK29) was screened for by BglII/PstI double digestion. Clones fitting this description were sequenced using

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

primers 108 and 109, which flank the insertion site in pWSK29, by SeqWright (Houston, TX, USA). An accurate clone was designated pSPN30.

Preparation of concentrated *S. Typhimurium* culture supernatant containing recombinant flagellins

S. Typhimurium strains were grown for 4 to 5 hours at 37°C with vigorous shaking by diluting an overnight culture 1 to 100 in 20 ml LB broth plus 1 mM IPTG. Once the OD₆₀₀ reached 0.8 to 1.2 bacteria were removed by centrifugation at 4000 rpm for 15 min and 12 ml of the resulting supernatant was passed through a 0.45 µm filter and subject to concentration by using an Amicon Ultra-15 with cutoff of 5K (Millipore) followed by a wash with 10 ml PBS. Protein concentration was determined by DC protein assay (BioRad) and SDS-PAGE followed by Coomassie blue stain. The final protein concentration was adjusted to 1 mg/ml.

Generation of rabbit anti-BaFliC serum and Western blot

B. abortus fliC (BaFliC) was amplified using primers BaFliC-F and BaFliC-R and cloned into pET103 in frame with a 6xHis tag. The resulting BaFliC::6xHis fusion protein was produced and purified by using Ni-NTA kit (Qiagen). Rabbit serum against BaFliC was generated by Antagene (Antagene Inc., Calif.). For detection of secreted BaFliC the supernatant from 1 ml culture was precipitated using trichloroacetic acid (TCA) and separated on a 12% SDS-PAGE gel. Proteins were electrotransferred to a polyvinylidene difluoride (PVDF) membrane. BaFliC was detected by using rabbit anti-BaFliC as primary antibody and as goat anti-rabbit IgG conjugated to horseradish peroxidase (HRP) as secondary antibody. *S. Typhimurium* Phase I flagellin (FliC) was detected using *Salmonella* Hi antiserum (Difco). C-terminal FLAG-tagged *S. Typhimurium* and *B. abortus* flagellins were detected using anti-FLAG monoclonal antibody (1:5000, Sigma) and a goat anti-mouse IgG antibody conjugated to

HRP. HRP activity was detected with a chemiluminescent substrate (PerkinElmer Life Sciences). Flagellin produced by *B. melitensis* 16M was detected as described previously (Fretin *et al.*, 2005).

Measurement of TLR5 agonist activity of flagellins

The human colonic epithelial cell line T-84 was cultured in were maintained in Dulbecco's modified Eagle medium (DMEM)-F12 medium (Gibco), containing 1.2 g/liter sodium bicarbonate, 2.5 mM L-glutamine, 15 mM HEPES, and 0.5 mM sodium pyruvate (Gibco), supplemented with 10% fetal calf serum (FCS). The day before assay cells from 1/3 of a 80 to 90% confluent T75 flask were seeded per each 24-well plate containing DMEM-F12 and 2% FCS. HEK293 cells were cultured as previously described (Kestra *et al.*, 2010). T84 cells in 24-well plates were either infected with 10 µl of bacteria grown as above or treated by adding 30 µl of concentrated bacterial culture supernatant and incubated for 4 hours at 37°C under 5% CO₂. For the HEK293 stably transfected with human TLR5, cells were grown in 48-well tissue culture plates and infected for 4-48 h with 10 µl of bacteria grown as described above or treated by adding 10 µl of concentrated bacteria culture supernatant and incubated for 8 hours at 37°C under 5% CO₂. Supernatants were aspirated and centrifuged for 10 min at 6,000 rpm to remove residual bacteria and cell debris before measurement of IL-8 concentration by ELISA.

Mitogen-activated protein kinase (MAPK) phosphorylation assay

T84 cells were seeded in six well plates at a density of 4×10^8 cells per well and incubated for 24h in DMEM/F12 + 10% FBS. The following day, cells were rinsed with PBS and the medium replaced with serum-free medium. For analysis of MAP kinase phosphorylation, cells were treated with concentrations of GST-BaFliC or GST-StFliC ranging from 250ng/ml

1
2
3 583 to 1 µg/ml. As a negative control, cells were treated with the highest concentration of flagellin
4
5 584 (1µg/ml) that had previously been treated with proteinase K (20mg/ml proteinase K for 1h at
6
7 585 37°C, then for 10 min at 75°C to inactivate the protease). After 30 and 90 min, cells were
8
9 586 lysed 0.1 ml in phosphosafe extraction reagent (Novagen) containing 2.5% protease inhibitor
10
11 587 (Sigma) according to the instructions of the manufacturer. The protein concentration was
12
13 588 determined using the Micro BCA kit (Pierce). Total protein (0.01 mg) was resolved by SDS-
14
15 589 PAGE and transferred to a polyvinylidene fluoride membrane. Primary antibodies were
16
17 590 purchased from Cell Signalling Technology, including the following phosphorylation-specific
18
19 591 antibodies: p-ERK and p-p38 (Thr180/Tyr182). Secondary antibodies (goat anti rabbit
20
21 592 conjugated to horseradish peroxidase) were purchased from Jackson Immunoresearch and
22
23 593 used according to the recommendations of the manufacturer. Peroxidase activity was
24
25 594 visualized using Immobilon Western Chemiluminescent HRP Substrate (Millipore). For each
26
27 595 primary antibody, a separate membrane was used.
28
29
30
31

32 596

33
34 597 **Detection of flagellin in the cytosol of infected macrophages**

35
36 598 The β-lactamase translocation assay was performed as previously described (Sun *et al.*, 2007).
37
38 599 Briefly J774A.1 mouse macrophages were seeded in 96-well coverglass bottom plates and
39
40 600 infected with *B. abortus* 2308 expressing either a BaFliC::Flag-TEM-1 fusion proteins, or an
41
42 601 irrelevant control (Glutathione-S-transferase::Flag-TEM-1) at a multiplicity of infection of
43
44 602 500. Plates were centrifuged at 250 g for 5 min at room temperature to synchronize infection.
45
46 603 After incubation for 1 hour at 37 °C in 5% CO₂, free bacteria were removed from the cells by
47
48 604 three washes with PBS. A volume of 0.2 ml of Dulbecco's modified Eagle's medium
49
50 605 supplemented with 10% heat-inactivated fetal bovine serum, 1% non-essential amino acids, 1
51
52 606 mM glutamine containing 1 mM IPTG and 100 ug/ml gentamicin was added to each well, and
53
54 607 plates were incubated at 37 °C in 5% CO₂. After 16 h, cells were washed once with Hank's
55
56
57
58
59
60

balanced salt solution (Invitrogen) and loaded with the fluorescent substrate CCF2/AM (1mM, Invitrogen) for 1.5 h at room temperature using the standard loading protocol recommended by the manufacturer. Fluorescence microscopy analysis was performed using an Axiovert M200 (Carl Zeiss), equipped with a CCF2 filter set (Chroma Technology). Fluorescence micrographs were captured using a Zeiss AxioCam MRC5 and Zeiss AxioVision 4.5 software. Images were imported into Adobe PhotoShop for color adjustment. The number of blue cells containing cleaved CCF2/AM was counted visually and expressed as the percentage of total cells in the well. The experiment was performed four times and the result expressed as geometric mean and range of the four experiments.

Bone-marrow derived Macrophages

Bone marrow-derived macrophages were isolated from C57BL/6, or congenic mutant mice following standard protocols as described previously (Sun *et al.*, 2007).

Macrophage infection

For assaying inflammasome activation, 24-well microtiter plates were seeded with bone marrow-derived macrophages at a concentration of 2×10^5 cells/well in 0.5 ml of RPMIsup and incubated over night at 37°C in 5% CO₂. For priming of macrophages, cells were treated for 4h before infection with LPS (100 ng/ml), as previously described (Franchi *et al.*, 2006). Inocula of *B. melitensis* 16M were prepared by growing with shaking in TSB for 24h. Bacteria were treated with a non-agglutinating (1:4,000) dilution of anti-*Brucella* rabbit serum (Difco) for 1h at 37 °C, as described (Rolan *et al.*, 2007) then diluted in RPMIsup to a concentration of 4×10^7 CFU/ml. Approximately 2×10^7 bacteria in 0.5 ml of RPMIsup, containing *B. melitensis* 16M wt or its isogenic *fliC* mutant, were added to each well of macrophages. Three independent assays were performed with triplicate samples, and each

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

experiment included control (C57BL/6) macrophages together with macrophages from mutant mice. Microtiter plates were centrifuged at 250 x g for 5 min at room temperature in order to synchronize infection. Cells were incubated for 20 min at 37°C in 5% CO₂, and free bacteria were removed by three washes with phosphate-buffered saline (PBS). RPMIsup plus 50mg gentamicin per ml was added to the wells, and the cells were incubated at 37°C in 5% CO₂. After 1 h, the RPMIsup plus 50µg/ml gentamicin was replaced with medium containing 25µg/ml gentamicin. Wells were sampled after infection by aspirating the medium, lysing the macrophages with 0.5 ml of 0.5% Tween-20 and rinsing each well with 0.5 ml of PBS. Viable bacteria were quantified by dilution in sterile PBS and plating on TSA containing appropriate antibiotics.

Liposome-mediated delivery of flagellins to the macrophage cytosol

Recombinant flagellin proteins were delivered to the macrophage cytosol using the cationic lipid DOTAP (Roche), as described previously (Franchi *et al.*, 2006). Briefly, 50 ml of DOTAP was incubated for 30 min in serum-free media with 2 mg of recombinant flagellins purified as described above. After incubation, 3.5 ml serum-free media was added and 500 µl was used to stimulate 1 x 10⁶ macrophages seeded in 24-well microtiter plates for 3h.

Measurement of cytokines

Mouse IL-1β was measured in culture supernatants by enzyme-linked immunoabsorbent assay (ELISA) (R&D Systems). Human IL-8 was detected using an ELISA kit from BioLegend.

Mice

1
2
3 657 Wild type (wt) BALB/c, wt C57BL/6, C57BL/6 *Nlrc4*^{-/-} (obtained from Dr. VM. Dixit and
4
5 658 described in (Mariathasan *et al.*, 2004)) and C57BL/6 *Casp1*^{-/-} (obtained from Dr. R. Flavell
6
7 659 and described in (Kuida *et al.*, 1995)) mice were used in this study. They were bred in the
8
9 660 animal facility of the University of Namur (Belgium). The animal handling and procedures of
10
11 661 this study were in accordance with the current European legislation (directive 86/609/EEC)
12
13 662 and in agreement with the corresponding Belgian law "*Arrêté royal relatif à la protection des*
14
15 663 *animaux d'expérience du 6 avril 2010 publié le 14 mai 2010*". The complete protocol was
16
17 664 reviewed and approved by the Animal Welfare Committee of the Facultés Universitaires
18
19 665 Notre-Dame de la Paix (FUNDP, Belgium)(Permit Number: 05-558).
20
21
22
23
24
25

26 667 **Infection of mice**

27
28 668 Mice were injected intraperitoneally (i.p.) with 4×10^4 CFUs of *B. melitensis* 16M in 500µl of
29
30 669 PBS. Control animals were injected with the same volume of PBS. Infectious doses were
31
32 670 validated by plating serial dilutions of the inocula. At selected time intervals, mice were
33
34 671 sacrificed by cervical dislocation. Immediately after being killed, spleen and liver were
35
36 672 collected for bacterial counts and histopathologic analyses. For bacterial counts, spleens and
37
38 673 livers were homogenized in PBS/0.1% X-100 triton (Sigma). Serial dilutions were plated on
39
40 674 2YT media plates for enumeration of tissue-associated CFU.
41
42
43
44
45

46 676 **Histology**

47
48 677 Spleens were fixed for 24h in Bouin's fixative, dehydrated for 24h in methanol, then
49
50 678 incubated in toluol and finally in warm paraffin prior to paraffin embedding. Sections (5µm)
51
52 679 were rehydrated and stained with hemalun, erythrosin and safran. Blinded histopathology
53
54 680 scoring for splenic granuloma formation was performed by a pathologist (MX), according to
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the following criteria. 0, <5% of splenic parenchyma containing granulomas; 1, 5-20%; 2, 20-40%; 3, 40-40%; 4, >60%.

Immunofluorescence microscopy

Spleens were fixed for 6h at 4°C in 2% paraformaldehyde (pH 7.4), washed in PBS, incubated overnight at 4°C in a 20% PBS-sucrose solution under agitation, and washed again in PBS. Tissues were embedded in the Tissue-Tek OCT compound (Sakura), frozen in liquid nitrogen, and cryostat sections (10µm) were prepared. Tissues sections were rehydrated in PBS, then incubated successively in a PBS solution containing 1% blocking reagent (Boeringer) (PBS-BR 1%) and in PBS-BR 1% containing any of the following mAbs or reagents: DAPI nucleic acid stain, Alexa Fluor 350 phalloidin, M1/70 (anti-CD11b, BD Biosciences), homemade anti-*B. melitensis* 16M serum (Copin *et al.*, 2012). Slides were mounted in Fluoro-Gel medium (Electron Microscopy Sciences, Hatfield, PA). Labelled tissues sections were visualized under a Zeiss fluorescent inverted microscope (Axiovert 200) equipped with high-resolution monochrome camera (AxioCam HR, Zeiss).

Statistical analysis

ANOVA I was used for data analysis after testing the homogeneity of variance (Bartlett test). Average comparisons were performed by pairwise Scheffe’s test. A Mann Whitney test was used for analysis of histopathology scoring. Errors bars represent standard deviation.

References

Akhter, A., Gavrilin, M.A., Frantz, L., Washington, S., Ditty, C., Limoli, D., *et al.* (2009). Caspase-7 activation by the Nlr4/Ipaf inflammasome restricts Legionella pneumophila infection. *PLoS Pathog* **5**, e1000361.

- 1
2
3 706 Amer, A., Franchi, L., Kanneganti, T.D., Body-Malapel, M., Ozoren, N., Brady, G., *et al.*
4
5 707 (2006). Regulation of Legionella phagosome maturation and infection through
6
7 708 flagellin and host Ipaf. *J Biol Chem* **281**, 35217-35223.
8
9
10 709 Andersen-Nissen, E., Smith, K.D., Strobe, K.L., Barrett, S.L., Cookson, B.T., Logan, S.M.
11
12 710 and Aderem, A. (2005). Evasion of Toll-like receptor 5 by flagellated bacteria. *Proc*
13
14 711 *Natl Acad Sci U S A* **102**, 9247-9252.
15
16 712 Anderson, T.D. and Cheville, N.F. (1986). Ultrastructural morphometric analysis of Brucella
17
18 713 abortus-infected trophoblasts in experimental placentitis. Bacterial replication occurs
19
20 714 in rough endoplasmic reticulum. *Am J Pathol* **124**, 226-237.
21
22
23 715 Arellano-Reynoso, B., Lapaque, N., Salcedo, S., Briones, G., Ciocchini, A.E., Ugalde, R., *et*
24
25 716 *al.* (2005). Cyclic beta-1,2-glucan is a Brucella virulence factor required for
26
27 717 intracellular survival. *Nat Immunol* **6**, 618-625.
28
29
30 718 Atluri, V.L., Xavier, M.N., de Jong, M.F., den Hartigh, A.B. and Tsolis, R.E. (2011).
31
32 719 Interactions of the human pathogenic Brucella species with their hosts. *Annu Rev*
33
34 720 *Microbiol* **65**, 523-541.
35
36 721 Ausubel, F.M., Brent, R., Kingston, R.E., E., M.D., Seidman, J.G., SMith, J.A. and Struhl, K.
37
38 722 (1991). Current protocols in Molecular Biology. *New-York: John Wiley & Sons.*
39
40
41 723 Barquero-Calvo, E., Chaves-Olarte, E., Weiss, D.S., Guzman-Verri, C., Chacon-Diaz, C.,
42
43 724 Rucavado, A., *et al.* (2007). Brucella abortus uses a stealthy strategy to avoid
44
45 725 activation of the innate immune system during the onset of infection. *PLoS One* **2**,
46
47 726 e631.
48
49
50 727 Beck, E., Ludwig, G., Auerswald, E.A., Reiss, B. and Schaller, H. (1982). Nucleotide
51
52 728 sequence and exact localization of the neomycin phosphotransferase gene from
53
54 729 transposon Tn5. *Gene* **19**, 327-336.
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Bergsbaken, T., Fink, S.L. and Cookson, B.T. (2009). Pyroptosis: host cell death and inflammation. *Nat Rev Microbiol* **7**, 99-109.

Briones, G., Inon de Iannino, N., Roset, M., Vigliocco, A., Paulo, P.S. and Ugalde, R.A. (2001). *Brucella abortus* cyclic beta-1,2-glucan mutants have reduced virulence in mice and are defective in intracellular replication in HeLa cells. *Infect Immun* **69**, 4528-4535.

Brodsky, I.E. and Medzhitov, R. (2009a). Targeting of immune signalling networks by bacterial pathogens. *Nat Cell Biol* **11**, 521-526.

Brodsky, I.E. and Monack, D. (2009b). NLR-mediated control of inflammasome assembly in the host response against bacterial pathogens. *Semin Immunol* **21**, 199-207.

Broz, P., Newton, K., Lamkanfi, M., Mariathasan, S., Dixit, V.M. and Monack, D.M. (2010). Redundant roles for inflammasome receptors NLRP3 and NLRC4 in host defense against *Salmonella*. *J Exp Med* **207**, 1745-1755.

Case, C.L., Shin, S. and Roy, C.R. (2009). Asc and Ipaf Inflammasomes direct distinct pathways for caspase-1 activation in response to *Legionella pneumophila*. *Infect Immun* **77**, 1981-1991.

Celli, J., de Chastellier, C., Franchini, D.M., Pizarro-Cerda, J., Moreno, E. and Gorvel, J.P. (2003). *Brucella* evades macrophage killing via VirB-dependent sustained interactions with the endoplasmic reticulum. *J Exp Med* **198**, 545-556.

Copin, R., De Baetselier, P., Carlier, Y., Letesson, J.J. and Muraille, E. (2007). MyD88-dependent activation of B220-CD11b+LY-6C+ dendritic cells during *Brucella melitensis* infection. *J Immunol* **178**, 5182-5191.

Copin, R., Vitry, M.A., Hanot Mambres, D., Machelart, A., De Trez, C., Vanderwinden, J.M., et al. (2012). In situ microscopy analysis reveals local innate immune response

- 754 developed around *Brucella* infected cells in resistant and susceptible mice. *PLoS*
755 *Pathog* **8**, e1002575.
- 756 Corbel, M.J. (1997). Brucellosis: an overview. *Emerg Infect Dis* **3**, 213-221.
- 757 Cummings, L.A., Wilkerson, W.D., Bergsbaken, T. and Cookson, B.T. (2006). In vivo, fliC
758 expression by *Salmonella enterica* serovar Typhimurium is heterogeneous, regulated
759 by ClpX, and anatomically restricted. *Mol Microbiol* **61**, 795-809.
- 760 Dinarello, C.A. (2009). Immunological and inflammatory functions of the interleukin-1
761 family. *Annu Rev Immunol* **27**, 519-550.
- 762 Eaves-Pyles, T., Murthy, K., Liaudet, L., Virag, L., Ross, G., Soriano, F.G., *et al.* (2001).
763 Flagellin, a novel mediator of *Salmonella*-induced epithelial activation and systemic
764 inflammation: I kappa B alpha degradation, induction of nitric oxide synthase,
765 induction of proinflammatory mediators, and cardiovascular dysfunction. *J Immunol*
766 **166**, 1248-1260.
- 767 Enright, F.M., Araya, L.N., Elzer, P.H., Rowe, G.E. and Winter, A.J. (1990). Comparative
768 histopathology in BALB/c mice infected with virulent and attenuated strains of
769 *Brucella abortus*. *Vet Immunol Immunopathol* **26**, 171-182.
- 770 Eskra, L., Canavessi, A., Carey, M. and Splitter, G. (2001). *Brucella abortus* genes identified
771 following constitutive growth and macrophage infection. *Infect Immun* **69**, 7736-7742.
- 772 Ferooz, J., Lemaire, J. and Letesson, J.J. (2011). Role of FlbT in flagellin production in
773 *Brucella melitensis*. *Microbiology* **157**, 1253-1262.
- 774 Feuillet, V., Medjane, S., Mondor, I., Demaria, O., Pagni, P.P., Galan, J.E., *et al.* (2006).
775 Involvement of Toll-like receptor 5 in the recognition of flagellated bacteria. *Proc*
776 *Natl Acad Sci U S A* **103**, 12487-12492.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Franchi, L., Amer, A., Body-Malapel, M., Kanneganti, T.D., Ozoren, N., Jagirdar, R., *et al.* (2006). Cytosolic flagellin requires Ipaf for activation of caspase-1 and interleukin 1beta in salmonella-infected macrophages. *Nat Immunol* **7**, 576-582.

Franchi L, Kamada N, Nakamura Y, Burberry A, Kuffa P, Suzuki S, Shaw MH, Kim YG, Núñez G. (2012) NLRC4-driven production of IL-1β discriminates between pathogenic and commensal bacteria and promotes host intestinal defense. *Nat Immunol.* **13**, 449-56.

Fretin, D., Fauconnier, A., Kohler, S., Halling, S., Leonard, S., Nijskens, C., *et al.* (2005). The sheathed flagellum of *Brucella melitensis* is involved in persistence in a murine model of infection. *Cell Microbiol* **7**, 687-698.

Gewirtz, A.T., Navas, T.A., Lyons, S., Godowski, P.J. and Madara, J.L. (2001). Cutting edge: bacterial flagellin activates basolaterally expressed TLR5 to induce epithelial proinflammatory gene expression. *J Immunol* **167**, 1882-1885.

Gross, A., Terraza, A., Ouahrani-Bettache, S., Liautard, J.P. and Dornand, J. (2000). In vitro *Brucella suis* infection prevents the programmed cell death of human monocytic cells. *Infect Immun* **68**, 342-351.

Grundling, A., Burrack, L.S., Bouwer, H.G. and Higgins, D.E. (2004). *Listeria monocytogenes* regulates flagellar motility gene expression through MogR, a transcriptional repressor required for virulence. *Proc Natl Acad Sci U S A* **101**, 12318-12323.

Hawn, T.R., Verbon, A., Lettinga, K.D., Zhao, L.P., Li, S.S., Laws, R.J., *et al.* (2003). A common dominant TLR5 stop codon polymorphism abolishes flagellin signaling and is associated with susceptibility to legionnaires' disease. *J Exp Med* **198**, 1563-1572.

- 800 Hayashi, F., Smith, K.D., Ozinsky, A., Hawn, T.R., Yi, E.C., Goodlett, D.R., *et al.* (2001).
801 The innate immune response to bacterial flagellin is mediated by Toll-like receptor 5.
802 *Nature* **410**, 1099-1103.
- 803 Hunt, A.C. and Bothwell, P.W. (1967). Histological findings in human brucellosis. *Journal of*
804 *clinical pathology* **20**, 267-272.
- 805 Kawai, T. and Akira, S. (2011). Toll-like Receptors and Their Crosstalk with Other Innate
806 Receptors in Infection and Immunity. *Immunity* **34**, 637-650.
- 807 Kestra, A.M., de Zoete, M.R., Bouwman, L.I. and van Putten, J.P. (2010). Chicken TLR21 is
808 an innate CpG DNA receptor distinct from mammalian TLR9. *J Immunol* **185**, 460-
809 467.
- 810 Kofoed, E.M. and Vance, R.E. (2011). Innate immune recognition of bacterial ligands by
811 NAIPs determines inflammasome specificity. *Nature* **477**, 592-595.
- 812 Kovach, M.E., Phillips, R.W., Elzer, P.H., Roop, R.M., 2nd and Peterson, K.M. (1994).
813 pBBR1MCS: a broad-host-range cloning vector. *BioTechniques* **16**, 800-802.
- 814 Kuida, K., Lippke, J.A., Ku, G., Harding, M.W., Livingston, D.J., Su, M.S. and Flavell, R.A.
815 (1995). Altered cytokine export and apoptosis in mice deficient in interleukin-1 beta
816 converting enzyme. *Science* **267**, 2000-2003.
- 817 Lamkanfi, M. and Dixit, V.M. (2009). Inflammasomes: guardians of cytosolic sanctity.
818 *Immunol Rev* **227**, 95-105.
- 819 Lapaque, N., Forquet, F., de Chastellier, C., Mishal, Z., Jolly, G., Moreno, E., *et al.* (2006).
820 Characterization of *Brucella abortus* lipopolysaccharide macrodomains as mega rafts.
821 *Cell Microbiol* **8**, 197-206.
- 822 Lapaque, N., Muller, A., Alexopoulou, L., Howard, J.C. and Gorvel, J.P. (2009). *Brucella*
823 *abortus* induces Irgm3 and Irga6 expression via type-I IFN by a MyD88-dependent

1
2
3 824 pathway, without the requirement of TLR2, TLR4, TLR5 and TLR9. *Microb Pathog*
4
5 825 **47**, 299-304.
6
7 826 Li, X., Lin, H., Zhang, W., Zou, Y., Zhang, J., Tang, X. and Zhou, J.M. (2005). Flagellin
8
9 827 induces innate immunity in nonhost interactions that is suppressed by *Pseudomonas*
10
11 828 *syringae* effectors. *Proc Natl Acad Sci U S A* **102**, 12990-12995.
12
13 829 Lightfield, K.L., Persson, J., Brubaker, S.W., Witte, C.E., von Moltke, J., Dunipace, E.A., *et*
14
15 830 *al.* (2008). Critical function for Naip5 in inflammasome activation by a conserved
16
17 831 carboxy-terminal domain of flagellin. *Nat Immunol* **9**, 1171-1178.
18
19 832 Lilleengen, K. (1948). Typing of *Salmonella typhimurium* by means of bacteriophage. *Acta.*
20
21 833 *Pathol. Microbiol. Scand. Suppl.* **77**, 2-125.
22
23 834 Mariathasan, S., Newton, K., Monack, D.M., Vucic, D., French, D.M., Lee, W.P., *et al.*
24
25 835 (2004). Differential activation of the inflammasome by caspase-1 adaptors ASC and
26
27 836 Ipaf. *Nature* **430**, 213-218.
28
29 837 McElvania Tekippe, E., Allen, I.C., Hulseberg, P.D., Sullivan, J.T., McCann, J.R., Sandor, M.,
30
31 838 *et al.* (2010). Granuloma formation and host defense in chronic *Mycobacterium*
32
33 839 tuberculosis infection requires PYCARD/ASC but not NLRP3 or caspase-1. *PLoS*
34
35 840 *One* **5**, e12320.
36
37 841 Miao, E.A., Alpuche-Aranda, C.M., Dors, M., Clark, A.E., Bader, M.W., Miller, S.I. and
38
39 842 Aderem, A. (2006). Cytoplasmic flagellin activates caspase-1 and secretion of
40
41 843 interleukin 1beta via Ipaf. *Nat Immunol* **7**, 569-575.
42
43 844 Miao, E.A., Andersen-Nissen, E., Warren, S.E. and Aderem, A. (2007). TLR5 and Ipaf: dual
44
45 845 sensors of bacterial flagellin in the innate immune system. *Semin Immunopathol* **29**,
46
47 846 275-288.
48
49
50
51
52
53
54
55
56
57
58
59
60

- 847 Miao, E.A., Leaf, I.A., Treuting, P.M., Mao, D.P., Dors, M., Sarkar, A., *et al.* (2010a).
848 Caspase-1-induced pyroptosis is an innate immune effector mechanism against
849 intracellular bacteria. *Nat Immunol* **11**, 1136-1142.
- 850 Miao, E.A., Mao, D.P., Yudkovsky, N., Bonneau, R., Lorang, C.G., Warren, S.E., *et al.*
851 (2010b). Innate immune detection of the type III secretion apparatus through the
852 NLRC4 inflammasome. *Proc Natl Acad Sci U S A* **107**, 3076-3080.
- 853 Molofsky, A.B., Byrne, B.G., Whitfield, N.N., Madigan, C.A., Fuse, E.T., Tateda, K. and
854 Swanson, M.S. (2006). Cytosolic recognition of flagellin by mouse macrophages
855 restricts *Legionella pneumophila* infection. *J Exp Med* **203**, 1093-1104.
- 856 Murphy, E.A., Sathiyaseelan, J., Parent, M.A., Zou, B. and Baldwin, C.L. (2001). Interferon-
857 gamma is crucial for surviving a *Brucella abortus* infection in both resistant C57BL/6
858 and susceptible BALB/c mice. *Immunology* **103**, 511-518.
- 859 O'Callaghan, D., Cazevaille, C., Allardet-Servent, A., Boschirolì, M.L., Bourg, G.,
860 Foulongne, V., *et al.* (1999). A homologue of the *Agrobacterium tumefaciens* VirB
861 and *Bordetella pertussis* Ptl type IV secretion systems is essential for intracellular
862 survival of *Brucella suis*. *Mol Microbiol* **33**, 1210-1220.
- 863 Pappas, G., Papadimitriou, P., Akritidis, N., Christou, L. and Tsianos, E.V. (2006). The new
864 global map of human brucellosis. *Lancet Infect Dis* **6**, 91-99.
- 865 Radhakrishnan, G.K., Yu, Q., Harms, J.S. and Splitter, G.A. (2009). *Brucella* TIR Domain-
866 containing Protein Mimics Properties of the Toll-like Receptor Adaptor Protein
867 TIRAP. *J Biol Chem* **284**, 9892-9898.
- 868 Raffatellu, M., Chessa, D., Wilson, R.P., Dusold, R., Rubino, S. and Baumler, A.J. (2005).
869 The Vi capsular antigen of *Salmonella enterica* serotype Typhi reduces Toll-like
870 receptor-dependent interleukin-8 expression in the intestinal mucosa. *Infect Immun* **73**,
871 3367-3374.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

872 Raupach, B., Peuschel, S.K., Monack, D.M. and Zychlinsky, A. (2006). Caspase-1-mediated
873 activation of interleukin-1beta (IL-1beta) and IL-18 contributes to innate immune
874 defenses against Salmonella enterica serovar Typhimurium infection. *Infect Immun* **74**,
875 4922-4926.

876 Ren, T., Zamboni, D.S., Roy, C.R., Dietrich, W.F. and Vance, R.E. (2006). Flagellin-deficient
877 Legionella mutants evade caspase-1- and Naip5-mediated macrophage immunity.
878 *PLoS Pathog* **2**, e18.

879 Rolan, H.G. and Tsolis, R.M. (2007). Mice lacking components of adaptive immunity show
880 increased Brucella abortus virB mutant colonization. *Infect Immun* **75**, 2965-2973.

881 Rolan, H.G., Xavier, M.N., Santos, R.L. and Tsolis, R.M. (2009). Natural antibody
882 contributes to host defense against an attenuated Brucella abortus virB mutant. *Infect*
883 *Immun* **77**, 3004-3013.

884 Salazar-Gonzalez, R.M. and McSorley, S.J. (2005). Salmonella flagellin, a microbial target of
885 the innate and adaptive immune system. *Immunol Lett* **101**, 117-122.

886 Salazar-Gonzalez, R.M., Srinivasan, A., Griffin, A., Muralimohan, G., Ertelt, J.M., Ravindran,
887 R., *et al.* (2007). Salmonella flagellin induces bystander activation of splenic dendritic
888 cells and hinders bacterial replication in vivo. *J Immunol* **179**, 6169-6175.

889 Salcedo, S.P., Marchesini, M.I., Lelouard, H., Fugier, E., Jolly, G., Balor, S., *et al.* (2008).
890 Brucella control of dendritic cell maturation is dependent on the TIR-containing
891 protein Btp1. *PLoS Pathog* **4**, e21.

892 Schroder, K. and Tschopp, J. (2010). The inflammasomes. *Cell* **140**, 821-832.

893 Sengupta, D., Koblansky, A., Gaines, J., Brown, T., West, A.P., Zhang, D., *et al.* (2009).
894 Subversion of innate immune responses by Brucella through the targeted degradation
895 of the TLR signaling adapter, MAL. *J Immunol* **184**, 956-964.

- 896 Shames, S.R. and Finlay, B.B. (2010). Breaking the stereotype: virulence factor-mediated
897 protection of host cells in bacterial pathogenesis. *PLoS Pathog* **6**, e1001057.
- 898 Simon, R., Priefer, U. and Puhler, A. (1983). A broad host range mobilization system for in
899 vivo genetic engineering: transposon mutagenesis in Gram-negative bacteria.
900 *Bio/Technology* **1**, 784-791.
- 901 Spink, W.W., Hoffbauer, F.W. and et al. (1949). Histopathology of the liver in human
902 brucellosis. *J Lab Clin Med* **34**, 40-58.
- 903 Stojiljkovic, I., Baumler, A.J. and Heffron, F. (1995). Ethanolamine utilization in *Salmonella*
904 typhimurium: nucleotide sequence, protein expression, and mutational analysis of the
905 cchA cchB eutE eutJ eutG eutH gene cluster. *Journal of bacteriology* **177**, 1357-1366.
- 906 Sun, Y.H., Rolan, H.G. and Tsolis, R.M. (2007). Injection of flagellin into the host cell
907 cytosol by *Salmonella enterica* serotype Typhimurium. *J Biol Chem* **282**, 33897-
908 33901.
- 909 van de Veerdonk, F.L., Netea, M.G., Dinarello, C.A. and Joosten, L.A. (2011).
910 Inflammasome activation and IL-1beta and IL-18 processing during infection. *Trends*
911 *Immunol* **32**, 110-116.
- 912 Vijay-Kumar, M., Wu, H., Jones, R., Grant, G., Babbitt, B., King, T.P., et al. (2006). Flagellin
913 suppresses epithelial apoptosis and limits disease during enteric infection. *Am J Pathol*
914 **169**, 1686-1700.
- 915 Wang, R.F. and Kushner, S.R. (1991). Construction of versatile low-copy-number vectors for
916 cloning, sequencing and gene expression in *Escherichia coli*. *Gene* **100**, 195-199.
- 917 Winter, S.E., Thiennimitr, P., Nuccio, S.P., Haneda, T., Winter, M.G., Wilson, R.P., et al.
918 (2009). Contribution of flagellin pattern recognition to intestinal inflammation during
919 *Salmonella enterica* serotype typhimurium infection. *Infect Immun* **77**, 1904-1916.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

920 Woodcock, D.M., Crowther, P.J., Doherty, J., Jefferson, S., DeCruz, E., Noyer-Weidner, M.,
921 *et al.* (1989). Quantitative evaluation of Escherichia coli host strains for tolerance to
922 cytosine methylation in plasmid and phage recombinants. *Nucleic acids research* **17**,
923 3469-3478.

924 Yonekura, K., Maki-Yonekura, S. and Namba, K. (2003). Complete atomic model of the
925 bacterial flagellar filament by electron cryomicroscopy. *Nature* **424**, 643-650.

926 Young, E.J. (1995). An overview of human brucellosis. *Clin Infect Dis* **21**, 283-289; quiz 290.

927 Yu, Y., Zeng, H., Lyons, S., Carlson, A., Merlin, D., Neish, A.S. and Gewirtz, A.T. (2003).
928 TLR5-mediated activation of p38 MAPK regulates epithelial IL-8 expression via
929 posttranscriptional mechanism. *Am J Physiol Gastrointest Liver Physiol* **285**, G282-
930 290.

931 Zamboni, D.S., Kobayashi, K.S., Kohlsdorf, T., Ogura, Y., Long, E.M., Vance, R.E., *et al.*
932 (2006). The Birc1e cytosolic pattern-recognition receptor contributes to the detection
933 and control of Legionella pneumophila infection. *Nat Immunol* **7**, 318-325.

934 Zhao, Y., Yang, J., Shi, J., Gong, Y.N., Lu, Q., Xu, H., *et al.* (2011). The NLRC4
935 inflammasome receptors for bacterial flagellin and type III secretion apparatus. *Nature*
936 **477**, 596-600.

937 Zygmunt, M.S., Hagius, S.D., Walker, J.V. and Elzer, P.H. (2006). Identification of Brucella
938 melitensis 16M genes required for bacterial survival in the caprine host. *Microbes*
939 *Infect* **8**, 2849-2854.

942 **Table 1. Bacterial strains and plasmids used in this study.**

<u>Designation</u>	<u>Genotype and/or Phenotype</u>	<u>Source or Reference</u>
Strains		
<i>Brucella melitensis</i> strains		
16M	wild type isolate	(Ferooz <i>et al.</i> , 2011)
$\Delta fliC$	$\Delta fliC::Kan$	
$\Delta flbT$	$\Delta flbT::Kan$	
$\Delta fliF$	$\Delta fliF::Kan$	
<i>BruFliC</i> ^{ON}	pBBR1- <i>fliC</i>	This work
<i>Brucella abortus</i> strain		
2308	wild type isolate	
<i>Salmonella enterica</i> serovar Typhimurium strains		
14028	ATCC 14028 Wild-Type	ATCC
IR715	14028 Spontaneous Nal ^R	
LT2	LT2 Wild-Type	(Lilleengen, 1948)
EHW26	IR715 <i>fliC::Tn10 fljB::MudJ (fliCfljB)</i>	(Raffatellu <i>et al.</i> , 2005)
<i>Escherichia coli</i> strains		
CC118 λ pir	<i>araD139</i> $\Delta(ara, leu)7697$ $\Delta lacX74$ <i>phoA</i> Δ 20 <i>galE galK thi rpsE rpoB argE_{am} recA1</i> λ pir	(Simon <i>et al.</i> , 1983)
DH10B	F- <i>mcrA</i> $\Delta(mrr-hsdRMS-mcrBC)$ $\phi 80 lacZ \Delta M15$ $\Delta lacX74$ <i>recA1 araD139</i> $\Delta(ara, leu)7697$ <i>galU galK rpsL</i> (Str ^R) <i>endA1 nupG</i>	Invitrogen
DH5 α	F- <i>mcrA</i> $\Delta(mrr-hsdRMS-mcrBC)$ $\phi 80 lacZ \Delta M15$ $\Delta(lacZYA-argF)U169$ <i>deoR recA1 endA1 phoA supE44</i> λ - <i>thi-1</i> <i>gyrA96 relA1</i>	(Woodcock <i>et al.</i> , 1989)
S17-1 λ pir	<i>recA thi pro r_K⁻ m_K⁺ RP4:2-Tc:MuKm Tn7</i> \square pir	(Simon <i>et al.</i> , 1983)
Plasmids		
pCR2.1	TOPO cloning vector	Invitrogen
pUC-KIXX	pUC4::Tn5 KanR	(Beck <i>et al.</i> , 1982)
pBBR1MCS	<i>mob RK2, lacZα, Cm^R</i>	(Kovach <i>et al.</i> , 1994)
pRH001 <i>fliC</i>	pMR10 (Cm ^R , <i>B. melitensis</i> 16M <i>Pflic-fliC-tfliC</i>)	This work
pBBR1- <i>fliC</i>	pBBR1MCS(Cm ^R , <i>B. melitensis</i> 16M <i>fliC</i>)	This work
pBBRFlag	pBBR1MCS::3xFLAG	This work
pYHS1116	pBBRFlag::StFliC	This work
pYHS1073	pBBRFlag::BaFliC	This work
pWSK29	Carb ^R , pSC101 <i>ori</i>	(Wang <i>et al.</i> , 1991)

943

Table 2: Primers used in this work

Primer	Sequence	Restriction site	Application
BaFliC-F	<u>ACCATATG</u> GCTAGCATTCTTACAAACTCGTCG	NdeI	FLAG-tagged BaFliC and BaFliC::FT fusion protein
BaFliC-R	<u>ACTGCAG</u> TTAGCCGCGGAACAGCGACAGGATCGAC	Sall	
StFliC-F	<u>ACCATATG</u> GCACAAGTCATTAATACAAACAGC	NdeI	FLAG-tagged StFliC and StFliC::FT fusion protein
StFliC-R	<u>AGTCGAC</u> TTAACGCAGTAAAGAGAGGACGTTT TGC	Sall	
StFliC-F2	<u>GAATTC</u> ATGGCACAAGTCATTAATACAAACAG C	EcoRI	GST-StFliC fusion protein
StFliC-STOP-R	<u>ACTCGA</u> GTTAACGCAGTAAAGAGAGGACGTTT TGC	XhoI	
BaFliC-F2	<u>GAATTC</u> ATGGCTAGCATTCTTACAAACTCG	EcoRI	GST fusion proteins
BaFliC-R2	<u>ACTCGAG</u> TTAGCCGCGGAACAGCGACAG	XhoI	GST fusion proteins
PfliC	<u>CGGGATCC</u> AATGCCCCGGGATCATGTTGATGC	BamHI	complementa tion plasmid
tfliC	<u>GCTCTAGAT</u> GCCAGACAGGATGTCGGGC	XbaI	
Plac	<u>GCTCtagA</u> tagAtagAGCGCAACGCAATTAATGT GAG	XbaI	fliC overexpressi on plasmid
fliC-Plac	GTTTGTAAGAATGCTAGCCATAGCTGTTTCCT GTGTGAAATTG		
BmfliC-F	ATGGCTAGCATTCTTACAAACTCGT		
BmfliC-R	<u>CGGGATCC</u> TTAGCCGCGGAACAGCG	BamHI	

Bold: Extra 5' DNA; **Bold/Underlined**: Multiple cloning site; ***Bold/Underlined/Italicized***:

Restriction site utilized in cloning; Lower case: Start or stop codon.

*

Figure legends

Fig. 1. Flagellin-deficient *B. melitensis* mutants infect macrophages *in vitro* with the same kinetics as wt bacteria but show enhanced persistence in mice.

(A) Western blot analysis of the production of flagellin (FliC, upper panel) by *B. melitensis* strains harvested at the early log phase and the log phase of growth in 2YT rich medium. Anti-Omp89 detection was used as a loading control (lower panel). Data are representative of two independent experiments. $\Delta fliC$ *pfliC* is the complemented strain. (B) Intracellular replication of *B. melitensis* 16M wt and $\Delta fliC$ strains in RAW264.7 murine macrophages. Error bars represent the standard deviation of triplicates in one representative experiment out of three. (C) Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated intraperitoneally (i.p.) with 4×10^4 CFUs of *B. melitensis* 16M wt, $\Delta fliC$, complemented $\Delta fliC$ *pfliC* (D) Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated intraperitoneally (i.p.) with 4×10^4 CFUs of *B. melitensis* 16M wt, $\Delta fliB$, or $\Delta fliF$ strains. Data represent the mean CFUs per organ and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). ** and *** denote highly significant ($p < 0.01$ and $p < 0.001$ respectively) differences in relation to wt infection. These results are representative of at least two independent experiments.

Fig. 2. Enhanced persistence of *B. melitensis* $\Delta fliC$ in mice is associated with increased pathology.

(A) Kinetics of splenomegaly in wt female BALB/c mice (n=5) injected i.p. with 4×10^4 CFUs of wt or $\Delta fliC$ strains of *B. melitensis* 16M. Data represent the mean spleen weight and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). *** denotes highly significant ($p < 0.001$) differences

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

in relation to wt infection. **(B)** Splenic pathology caused by a 28 day-infection was determined using the histopathology scoring system as described in the Material and methods. Data were analysed using a Mann Whitney test, and the mean histopathology scores were significantly different ($P=0.009$) **(C)** Representative photomicrographs (x10) of histopathology of spleens from BALB/c mice uninfected or infected for 28 days with *B. melitensis* wt or $\Delta fliC$ strain. WP, white pulp; T, thrombosis; black arrows, granuloma; white arrowhead, neutrophil infiltration. These results are representative of at least two independent experiments.

Fig. 3. Constitutive production of flagellin does not impair replication of *B. melitensis* 16M in macrophages *in vitro*, but attenuates its virulence *in vivo*.

(A) Western blot analysis of flagellin (FliC, upper panel) production in wt and *BruFliC*^{ON} strains during early exponential and stationary phases of growth in 2YT rich medium. Detection of Omp89 was used as a loading control. **(B)** Intracellular replication of wt and *BruFliC*^{ON} strains in RAW264.7 murine macrophages. Error bars represent the standard deviation of triplicates in one representative experiment out of two. **(C)** Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated i.p. with 4×10^4 CFUs of wt or *BruFliC*^{ON} strain. Data represent the mean CFUs per organ and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). ** and *** denote highly significant ($p<0.01$ and $p<0.001$ respectively) differences in relation to wt infection. These results are representative of at least two independent experiments.

Fig. 4. *Brucella* flagellin lacks TLR5 agonist activity.

(A-C) FLAG-tagged flagellins from *S. enterica* serotype Typhimurium (StFliC) or *Brucella abortus* (BaFliC) were expressed in an *S. Typhimurium fliCfljB* mutant, and culture

supernatants containing recombinant flagellins were used to treat cells. **(A)** Western blot showing production of bacterium-associated flagellins from *S. Typhimurium* wt (lane 1), *S. Typhimurium fliCfljB* mutant (lane 2), *fliCfljB* mutant expressing StFliC-FLAG (lane 3) or *fliCfljB* mutant expressing BaFliC-FLAG (lane 4). Flagellins were detected both in the pellets (left panel) and in the concentrated supernatants (right panel) of *S. Typhimurium* strains. 30ng of concentrated supernatant proteins from *S. Typhimurium* strains expressing recombinant flagellins were used to treat HEK293/hTLR5 cells for 4 or 24h **(B)** and T84 cells for 8h **(C)**. IL-8 in cell supernatants was measured by ELISA. **(D)** Activation of p38 and ERK MAPK in T-84 cells by purified recombinant flagellins from *Brucella* (GST-BaFliC) and *S. Typhimurium* (GST-StFliC) was measured by Western blot analysis with anti- p38, anti-phosphorylated (P-)p38, anti- ERK, and anti-P-ERK. Detection of tubulin was used as a loading control. Purified flagellins treated with proteinase K (PK) were used as a control. All data shown are from an individual experiment that was repeated at least twice with similar results.

Fig. 5. *B. abortus* flagellin can enter the cytosol of infected macrophages and induces IL-1 β in an NLRC4-independent manner. **(A)** Bone marrow-derived macrophages from C57BL/6 mice were inoculated with *B. melitensis* 16M wt or the $\Delta fliC$ mutant and IL-1 β was measured in the culture supernatants by ELISA at 24h p.i. Results are shown as the mean \pm standard deviation of data from five independent experiments. **(B)** Bone marrow-derived macrophages from C57BL/6 or *Nlrc4*^{-/-} mice were inoculated with *B. melitensis* 16M wt or the *BruFliC*^{ON} strain. IL-1 β in the supernatant was measured at 6h after inoculation. Data shown are combined from three independent experiments with triplicate samples, and represent the mean \pm standard deviation of all data.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Fig. 6. Introduction of recombinant *Brucella* flagellin into the host cell cytosol results in ASC-dependent, but NLRC4-independent secretion of IL-1 β .

Graded amounts of GST-BaFliC and GST-StFliC fusion proteins were delivered to the cytosol of LPS-primed bone marrow-derived macrophages from C57BL/6 (A), *Nlrc4*^{-/-} (B) or *Asc*^{-/-} (C) mice, using the cationic lipid DOTAP. Treated macrophages were incubated for 3h before measurement of IL-1 β in the supernatants by ELISA. Results are expressed as the mean of triplicate samples, with error bars representing the range of the data from one of two independent experiments with the same outcome.

Fig. 7. NLRC4 inflammasome is implicated in the control of *B. melitensis* infection in vivo. Wild type, *Nlrc4*^{-/-} (A) and *Casp1*^{-/-} (B) C57BL/6 mice (n=5) were injected i.p. with 4 x 10⁴ CFUs of *B. melitensis* wt, *BruFliC*^{ON} or Δ *fliC* strain, as indicated in the figure. Mice were sacrificed 21 days post-infection and CFUs per spleen were determined. These results are representative of at least two independent experiments. Data have been analysed by ANOVA I after testing the homogeneity of variance (Bartlett). * and ** denote respectively significant (p<0.05) and highly significant (p<0.01) differences in relation to C57BL/6 wt infection by wt bacteria.

Fig. 8. The distribution of *Bru*-positive cells is different in the spleen of mice infected by the Δ *fliC* mutant, compared to wt infection.

Localization of *Bru*⁺ cells (green) and CD11b⁺ cells (red) in the spleen of BALB/c mice non-infected or infected with *B. melitensis* wt or the Δ *fliC* strain. The graph represents the relative number of clusters of *Bru*⁺ cells. Errors bars are the standard deviation calculated on countings of four mice from two independent experiments.

1049 Acknowledgements

1050 We thank V. Dixit, J. Tschopp and A. Tardivel for providing us with the NLRC4 KO mice.

1051 Part of this work has been granted by an ARC Convention from the French community of

1052 Belgium (N° 08/13-015). M. Terwagne holds a PhD grant from FNRS (Fond National pour la

1053 Recherche Scientifique) and J. Ferooz holds a PhD grant from FRIA (Fonds pour la formation

1054 à la Recherche dans l'Industrie et l'Agriculture).

1055 This work was supported by US PHS grants AI50553 and AI097107 to R.M.T. and US PHS

1056 grant DK091191 to G.N. V.L.A was supported by T32 IA60555.

1057




Fig. 1. Flagellin-deficient *B. melitensis* mutants infect macrophages in vitro with the same kinetics as wt bacteria but show enhanced persistence in mice. (A) Western blot analysis of the production of flagellin (FliC, upper panel) by *B. melitensis* strains harvested at the early log phase and the log phase of growth in 2YT rich medium. Anti-Omp89 detection was used as a loading control (lower panel). Data are representative of two independent experiments. $\Delta fliC$ pflIC is the complemented strain. (B) Intracellular replication of *B. melitensis* 16M wt and $\Delta fliC$ strains in RAW264.7 murine macrophages. Error bars represent the standard deviation of triplicates in one representative experiment out of three. (C) Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated intraperitoneally (i.p.) with 4×10^4 CFUs of *B. melitensis* 16M wt, $\Delta fliC$, complemented $\Delta fliC$ pflIC (D) Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated intraperitoneally (i.p.) with 4×10^4 CFUs of *B. melitensis* 16M wt, $\Delta flbT$, or $\Delta fliF$ strains. Data represent the mean CFUs per organ and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). ** and *** denote highly significant ($p < 0.01$ and $p < 0.001$ respectively) differences in relation to wt infection. These results are representative of at least two independent experiments.

1057x793mm (72 x 72 DPI)




Fig. 2. Enhanced persistence of *B. melitensis* $\Delta fliC$ in mice is associated with increased pathology. (A) Kinetics of splenomegaly in wt female BALB/c mice ($n=5$) injected i.p. with 4×10^4 CFUs of wt or $\Delta fliC$ strains of *B. melitensis* 16M. Data represent the mean spleen weight and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). *** denotes highly significant ($p<0.001$) differences in relation to wt infection. (B) Splenic pathology caused by a 28 day-infection was determined using the histopathology scoring system as described in the Material and methods. Data were analysed using a Mann Whitney test, and the mean histopathology scores were significantly different ($P=0.009$) (C) Representative photomicrographs ($\times 10$) of histopathology of spleens from BALB/c mice uninfected or infected for 28 days with *B. melitensis* wt or $\Delta fliC$ strain. WP, white pulp; T, thrombosis; black arrows, granuloma; white arrowhead, neutrophil infiltration. These results are representative of at least two independent experiments.

793x595mm (72 x 72 DPI)




Fig. 3. Constitutive production of flagellin does not impair replication of *B. melitensis* 16M in macrophages in vitro, but attenuates its virulence in vivo.

(A) Western blot analysis of flagellin (FliC, upper panel) production in wt and *Bru*FliC^{ON} strains during early exponential and stationary phases of growth in 2YT rich medium. Detection of Omp89 was used as a loading control. (B) Intracellular replication of wt and *Bru*FliC^{ON} strains in RAW264.7 murine macrophages. Error bars represent the standard deviation of triplicates in one representative experiment out of two. (C) Infection kinetics in the spleens of wt BALB/c mice (n=5) inoculated i.p. with 4 x 10⁴ CFUs of wt or *Bru*FliC^{ON} strain. Data represent the mean CFUs per organ and error bars represent standard deviation. Results have been analyzed by ANOVA I after testing the homogeneity of variance (Bartlett). ** and *** denote highly significant (p<0.01 and p<0.001 respectively) differences in relation to wt infection. These results are representative of at least two independent experiments.

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Fig. 4. *Brucella* flagellin lacks TLR5 agonist activity.

(A-C) FLAG-tagged flagellins from *S. enterica* serotype Typhimurium (StFliC) or *Brucella abortus* (BaFliC) were expressed in an *S. Typhimurium* *fliCfljB* mutant, and culture supernatants containing recombinant flagellins were used to treat cells. (A) Western blot showing production of bacterium-associated flagellins from *S. Typhimurium* wt (lane 1), *S. Typhimurium* *fliCfljB*mutant (lane 2), *fliCfljB* mutant expressing StFliC-FLAG (lane 3) or *fliCfljB* mutant expressing BaFliC-FLAG (lane 4). Flagellins were detected both in the pellets (left panel) and in the concentrated supernatants (right panel) of *S. Typhimurium* strains. 30ng of concentrated supernatant proteins from *S. Typhimurium* strains expressing recombinant flagellins were used to treat HEK293/hTLR5 cells for 4 or 24h (B) and T84 cells for 8h (C). IL-8 in cell supernatants was measured by ELISA. (D) Activation of p38 and ERK MAPK in T-84 cells by purified recombinant flagellins from *Brucella* (GST-BaFliC) and *S. Typhimurium* (GST-StFliC) was measured by Western blot analysis with anti- p38, anti-phosphorylated (P-)p38, anti- ERK, and anti-P-ERK. Detection of tubulin was used as a loading control. Purified flagellins treated with proteinase K (PK) were used as a control. All data shown are

from an individual experiment that was repeated at least twice with similar results.

595x793mm (72 x 72 DPI)

For Peer Review




Fig. 5. *B. abortus* flagellin can enter the cytosol of infected macrophages and induces IL-1 β in an NLRC4-independent manner. (A) Bone marrow-derived macrophages from C57BL/6 mice were inoculated with *B. melitensis* 16M wt or the $\Delta fliC$ mutant and IL-1 β was measured in the culture supernatants by ELISA at 24h p.i. Results are shown as the mean \pm standard deviation of data from five independent experiments. (B) Bone marrow-derived macrophages from C57BL/6 or Nlrc4 $^{-/-}$ mice were inoculated with *B. melitensis* 16M wt or the BruFliCONstrain. IL-1 β in the supernatant was measured at 6h after inoculation. Data shown are combined from three independent experiments with triplicate samples, and represent the mean \pm standard deviation of all data.

595x793mm (72 x 72 DPI)




Fig. 6. Introduction of recombinant Brucella flagellin into the host cell cytosol results in ASC-dependent, but NLRC4-independent secretion of IL-1 β . Graded amounts of GST-BaFliC and GST-StFliC fusion proteins were delivered to the cytosol of LPS-primed bone marrow-derived macrophages from C57BL/6 (A), Nlrc4^{-/-} (B) or Asc^{-/-} (C) mice, using the cationic lipid DOTAP. Treated macrophages were incubated for 3h before measurement of IL-1 β in the supernatants by ELISA. Results are expressed as the mean of triplicate samples, with error bars representing the range of the data from one of two independent experiments with the same outcome.

595x793mm (72 x 72 DPI)




Fig. 7. NLRC4 inflammasome is implicated in the control of *B. melitensis* infection in vivo. Wild type, *Nlrc4*^{-/-} (A) and *Casp1*^{-/-} (B) C57BL/6 mice (n=5) were injected i.p. with 4 x 10⁴ CFUs of *B. melitensis* wt, *BruFliCON* or $\Delta fliC$ strain, as indicated in the figure. Mice were sacrificed 21 days post-infection and CFUs per spleen were determined. These results are representative of at least two independent experiments. Data have been analysed by ANOVA I after testing the homogeneity of variance (Bartlett). * and ** denote respectively significant (p<0.05) and highly significant (p<0.01) differences in relation to C57BL/6 wt infection by wt bacteria.




Fig. 8. The distribution of Bru-positive cells is different in the spleen of mice infected by the $\Delta fliC$ mutant, compared to wt infection.

Localization of Bru+ cells (green) and CD11b+ cells (red) in the spleen of BALB/c mice non-infected or infected with *B. melitensis* wt or the $\Delta fliC$ strain. The graph represents the relative number of clusters of Bru+ cells. Errors bars are the standard deviation calculated on countings of four mice from two independent experiments.

793x595mm (72 x 72 DPI)